

THE SECOND ALL **BREAST CANCER** ***REPORT***

Focussing on
Inequalities:
Variation in
breast cancer
outcomes with
age and deprivation



West Midlands
Cancer Intelligence Unit



Acknowledgements

The authors of this report:

| | |
|----------------------------|--|
| Dr Gill Lawrence | Director, West Midlands Cancer Intelligence Unit Regional Director of Breast Screening Quality Assurance |
| Ms Olive Kearins | Deputy Regional Director of Breast Screening Quality Assurance West Midlands Breast Screening QA Reference Centre |
| Dr Catherine Lagord | Breast Cancer Audit Project Manager West Midlands Cancer Intelligence Unit |
| Ms Shan Cheung | Breast Screening QA Senior Information Analyst West Midlands Breast Screening QA Reference Centre |
| Dr Jasmin Sidhu | Breast Cancer Data Analyst West Midlands Cancer Intelligence Unit |
| Mr Colin Sagar | BCCOM Information Assistant West Midlands Cancer Intelligence Unit |

would like to express their thanks to the following organisations and individuals for their contributions to the second *'All Breast Cancer Report'*;

UK cancer registries which provided the core 2007 data on all breast cancers diagnosed in their populations.

English regional breast screening QA reference centres which provided the identifiers to enable the screen-detected cancers originally submitted to the NHS Breast Screening Programme audit of screen-detected cancers to be matched to the cancer registry data.

Mr Alan MacDonald, Computer Information Systems Development Specialist at the West Midlands Cancer Intelligence Unit who undertook the matching of the various datasets.

Dr Yoon C Chia, Consultant Pathologist at Wycombe Hospital who provided pathology graphics for this report.

Breast surgeons across the UK who checked and submitted data into the NHS Breast Screening Programme / Association of Breast Surgery audit of screen-detected breast cancer and to the Breast Cancer Clinical Outcome Measures (BCCOM) audit (see Appendix 1)

The authors would also like to express their thanks to the National Cancer Intelligence Network for their financial assistance in the production of this report.



This is the second All Breast Cancer Report and draws information from a number of sources on over 50,000 people with newly diagnosed breast tumours in the UK in 2007, of whom 89% had invasive disease.

Despite a continuing overall improvement in UK breast cancer outcomes, significant

differences remain in the presentation and management of the disease that are associated with age, deprivation and ethnicity. This report focuses on the differences related to age and deprivation. It sends out clear messages on variations in treatment and differences in outcome and shows where we can focus our attention to address variations in stage at diagnosis and treatment provided.

There are some very positive messages on experience and outcome that demonstrate real improvements for patients: rates of breast conservation surgery are high,

immediate breast reconstruction is being performed more often for women undergoing mastectomy, and 1-year and 5-year relative survival rates are increasing. The beneficial impact of the NHS Breast Screening Programme is clearly apparent in the predominance of cancers that are small at diagnosis, and in its effect of reducing the inequalities associated with age and deprivation.

Whilst I very much welcome this report, and commend the detailed and dedicated work that has been put in to produce it, there is a concern that it is based on UK data from 2007. If we are to be more effective in the future in reducing variation and dealing with inequality, we will require information that is more contemporaneous, and it is therefore encouraging to know that a more streamlined approach to national cancer data collection is being developed that should enable this.

Mr Martin Lee

**Chair, National Cancer Intelligence Network
Breast Clinical Reference Group**

Contents

| | Page No. |
|---|-----------------|
| Executive Summary | 1 |
| Introduction and Data Sources | 3 |
| Introduction | 3 |
| Data Sources | 3 |
| Cohort Characteristics | 5 |
| Key Findings | 5 |
| Country Profile | 5 |
| Age, Gender and Deprivation | 6 |
| Route of Presentation | 8 |
| Ethnicity | 9 |
| Tumour Characteristics | 11 |
| Key Findings | 11 |
| Tumour Characteristics | 11 |
| Invasive Tumour Size | 11 |
| Invasive Tumour Grade | 12 |
| Nodal Status | 13 |
| Nottingham Prognostic Index (NPI) | 14 |
| Vascular Invasion | 15 |
| Oestrogen Receptor Status (ER Status) | 16 |
| Human Epidermal Growth Factor Receptor 2 Status (HER2 Status) | 17 |
| Surgical Treatment Outcomes | 19 |
| Key Findings | 19 |
| Surgical Treatment | 20 |
| Breast Conserving Surgery and Mastectomy | 21 |
| Mastectomy: Breast Conserving Surgery Ratio | 22 |
| Immediate Reconstruction | 23 |
| Sentinel Lymph Node Biopsy (SLNB) | 25 |
| Length of Hospital Stay | 26 |
| Repeat Operation Rates | 27 |
| 30 Day Mortality | 28 |
| Surgical Caseload | 29 |
| Survival | 30 |
| Key Findings | 30 |
| Relative Survival | 30 |
| Mortality | 31 |
| 1-year and 5-year Relative Survival | 31 |
| Age | 32 |
| Deprivation | 33 |
| References | 35 |
| Appendix 1 – Participating Surgeons | 36 |
| Appendix 2 – Overall Data Quality and Completeness | 39 |
| Appendix 3 – Tumour Characteristics Data Completeness | 41 |
| Appendix 4 – Acronyms | 42 |

Executive Summary

Introduction

There is an increasing need to publish data on the outcomes of care - what actually happens to the health of the patient as a result of the treatment and care they receive. In order to place a stronger focus on clinical issues, the *National Cancer Peer Review Evidence Guide* published in February 2010 introduced clinical lines of enquiry into the review process. In this, the second *All Breast Cancer Report*, as well as including chapters on cohort and tumour characteristics similar to those in the first *All Breast Cancer Report*, we have focussed on surgical treatment and how this varies with age, deprivation and route of presentation. The analyses

Cohort Characteristics

Of the 50,286 cases of invasive (44,782; 89%) and non/micro-invasive (5,490; 11%) breast cancer diagnosed in the UK in 2007, 292 (0.6%) were diagnosed in men. 81% were diagnosed in patients aged 50 and over, and 32% of all breast cancers and 56% of breast cancers diagnosed in women between the ages of 50 and 69

Tumour Characteristics

Of the surgically treated invasive breast cancers, 58% were small with an invasive tumour size of 20mm or less, 17% were grade 1, 62% were lymph node negative, 39% were in the excellent (EPG) and good (GPG) Nottingham Prognostic Index prognostic groups, 29% had vascular invasion, 84% were ER positive and 15% were HER2 positive.

Screen-detected invasive breast cancers were more likely to be smaller in size, of a lower grade, node negative and in the EPG and GPG prognostic groups. Screen-detected cancers were also less likely to have vascular invasion, more likely to be ER positive and less likely to be HER2 positive. These differences between symptomatic and screen-detected cancers were evident across all ages and deprivation quintiles.

include, for breast cancers diagnosed in women in 2007, the cancer peer review headline clinical indicators:

- access to immediate reconstruction
- ratio of mastectomy to breast conserving surgery
- surgical caseload
- average length of stay
- 1-year and 5-year relative survival rates.

The report also includes analyses on

- the use of sentinel lymph node biopsy
- repeat operations
- 30 day mortality.

years, were screen-detected. Breast cancer increases with affluence; 23% of cases were diagnosed in the least deprived group compared with 15% in the most deprived group. Three quarters of cases in English women were diagnosed in White women (20% unknown ethnicity).

Compared with women in the screening age group, older women were more likely to have node negative cancers, and less likely to have small cancers, HER2 positive cancers and vascular invasion present. Women aged less than 40 were less likely to have grade 1 tumours and cancers in the EPG and GPG prognostic groups and more likely to have vascular invasion present.

For symptomatic invasive breast cancers, the proportions with good prognosis (small, node negative, ER positive, EPG and GPG prognostic groups) decreased with increasing deprivation. Equivalent trends were not apparent for screen-detected cancers, indicating that screening reduces the observable inequalities in tumour characteristics associated with deprivation.

Executive Summary

Surgical Treatment

82% of women diagnosed with breast cancer in 2007 had surgery, and 57% of those had a breast conserving procedure. Significantly more women with screen-detected breast cancer had surgery (98% compared with 75% for those presenting symptomatically), and significantly fewer had a mastectomy (27% compared with 53%). Of the 524 surgeons who treated female breast cancer patients in 2007, 139 (27%) had a caseload of less than 30 patients, 27 had a significantly high mastectomy rate and 44 a significantly low mastectomy rate.

For symptomatic breast cancers, surgical treatment decreased with age at diagnosis; with only 74% of women aged 70-79 and 39% of women aged 80 and above having surgery compared with 90% of women aged under 50. Women in the most deprived quintile were less likely to have surgery than those in the least deprived quintile (72% compared with 78%). This effect was not apparent for women with screen-detected breast cancers.

Overall, 11% of female mastectomy patients had immediate reconstruction. A greater proportion of non/

micro-invasive cancers had immediate reconstruction (27% compared with 10% of invasive cancers), and women with screen-detected cancers were more likely to have an immediate reconstruction (16% versus 10%). For screen-detected cancers, immediate reconstruction was more likely to be undertaken in women from a less deprived background. 32% of women had a sentinel lymph node biopsy (SLNB). Older women were less likely to have a SLNB.

The median length of stay for female mastectomy patients was 4 days compared to 2 days for women having breast conserving surgery. Screening patients had a shorter length of stay, mainly because more were treated with breast conserving surgery. Repeat operation rates were higher for non/micro-invasive cancers (25% compared with 14% for invasive cancers), for women with screen-detected cancer (18%) and for younger women (19%). 860 women (2%) died within 30 days of their diagnosis. 51% of these were death certificate only (DCO) cases. Of the non-DCO cases who died within 30 days, 64% died of breast cancer. A significantly higher proportion of women in the most deprived quintile died within 30 days of their cancer diagnosis.

Survival

1-year relative survival was significantly higher for women diagnosed with breast cancer in 2007 (96% compared to 94% in 2002/03). 1-year and 5-year relative survival rates were significantly higher for women with screen-detected breast cancer (100% compared to 93%-94% at 1 year and 97% compared to 77% at 5 years). For women with symptomatic breast cancer, 5-year relative survival decreased with age (from 86% in women aged 40-49 years to only 62% in women aged 80 years and above).

For women with symptomatic breast cancer, there were marked decreases in 1-year survival with deprivation

(from 90%-92% for the most deprived women to 96%-97% for the least deprived women). These differences were not apparent for women with screen-detected breast cancer. Although 5-year relative survival for women with screen-detected cancer did decrease from 99% in the least deprived to 94% in the most deprived, there was a much more marked difference for women with symptomatic breast cancer; 5-year relative survival being 83% in the least deprived and 68% in the most deprived. Screening thus appears to reduce the observable inequalities in 1-year and 5-year relative survival associated with deprivation.

Introduction and Data Sources

Introduction

There is an increasing need to publish data on the outcomes of care - what actually happens to the health of the patient as a result of the treatment and care they receive. The White Paper *Equity and Excellence: Liberating the NHS*¹ sets out how improvements in outcomes will in future be the primary focus of the NHS. The associated document *Liberating the NHS: Transparency in outcomes; a framework for the NHS*² discusses how these outcomes will be measured. Breast cancer is, in some ways, ahead of the crowd in having clinical outcome measures defined by the Breast Cancer Clinical Outcome Measures (BCCOM) Project and an established track record of audit and reporting the processes of care at national level for screen-detected and symptomatic breast cancers. How breast cancer outcomes will be measured in future at national level will become clearer when the *National Institute for Health and Clinical Excellence (NICE) Quality Standards for Breast Cancer*³ are published later this year.

In order to place a stronger focus on clinical issues to make cancer peer review clinically relevant and to sustain the continued support and involvement of clinical staff, the *National Cancer Peer Review Evidence*

Guide published in February 2010⁴ introduced clinical lines of enquiry into the review process. As part of this initiative, a number of headline clinical indicators were agreed with the National Cancer Intelligence Network (NCIN) Breast Clinical Reference Group (CRG), some of which are based on data collected at a national level.

In this, the second *All Breast Cancer Report* as well as including chapters on cohort and tumour characteristics similar to those in the first *All Breast Cancer Report*⁵, we have focussed on surgical treatment and how this varies with age, deprivation status and route of presentation. The analyses include the cancer peer review headline clinical indicators:

- access to immediate reconstruction
- ratio of mastectomy to breast conserving surgery
- surgical caseload
- average length of stay
- 1-year and 5-year relative survival rates.

The report also includes analyses on

- the use of sentinel lymph node biopsy
- repeat operations
- 30 day mortality.

Data Sources

Data from the different sources were linked using NHS number and amalgamated to produce a single record for each patient.

Cancer registry data – Population based data on the diagnosis, treatment and survival of breast cancer cases in the United Kingdom (UK) are collected by cancer registries in eight English regions and in Northern Ireland, Scotland and Wales (the Celtic countries). The dataset collected is defined in the Cancer Registration Minimum Data Set⁶.

HES data – Hospital Episode Statistics (HES)⁷ record details of the care provided by NHS hospitals in England. Information on patient age, gender and ethnicity,

clinical diagnoses and operations, time waited and admission dates are recorded in HES. Details of self-reported ethnicity, surgery, chemotherapy and selected patient demographics from HES were used to supplement the cancer registry data. In Northern Ireland, hospital information is obtained from the Decision Support System through the cancer registry database. The Patient Episode Database for Wales (PEDW) records all episodes of inpatient or daycase activity in NHS Wales hospitals. The primary source of hospital level information in Scotland is the Scottish Morbidity Record (SMR01) which contains hospital inpatient or daycase discharge records.

Introduction and Data Sources

ONS data – The Office for National Statistics (ONS) collates data held by the English regional cancer registries and the Welsh Cancer Intelligence and Surveillance Unit to give aggregated data for England and Wales. The ONS cancer registration dataset contains patient demographics, some tumour characteristics and treatment flags indicating that the patient has had surgery, radiotherapy and/or chemotherapy. The ONS and HES datasets, were combined to obtain a database of all registered breast cancer patients, their demographics and their in-patient treatment.

Indices of Multiple Deprivation – The Index of Multiple Deprivation 2007 (ID2007) combines a number of indicators, covering a range of economic, social and housing issues, into a single deprivation score for each small area in England. This allows each area to be ranked according to its level of deprivation. ID2007 scores are produced at Lower Super Output Area (LSOA) level, of which there are 32,482 in England⁸. The income domain score was used as the deprivation indicator for England in this report. Income domain scores are grouped into 5 ranges (quintiles), each containing one fifth of the English population. To obtain an indication of the deprivation status of each breast cancer patient, postcode of residence was linked to the income domain score for the small area in which the patient lived at the time of diagnosis. Patients were then allocated to a deprivation quintile based on their score.

The Northern Ireland Cancer Registry uses the Output Area (OA) level Economic Deprivation Measure of the Northern Ireland Multiple Deprivation Measure (NIMDM) 2005 to derive deprivation scores. This measure combines income, employment and proximity to services domains, and used in conjunction with the

patient's postcode was put into deprivation quintiles. The Welsh Cancer Intelligence and Surveillance Unit uses the Welsh Index of Multiple Deprivation (WIMD) 2008 to derive deprivation scores. The LSOA of the patient's residence was used to assign a quintile based upon the income domain using equal population in each quintile. In Scotland, patient postcodes are utilised in the generation of the Scottish Index of Multiple Deprivation (SIMD) 2006 which is then used to derive deprivation scores.

BCCOM audit validated data – Each year, to initiate the Breast Cancer Clinical Outcome Measures (BCCOM) audit, data for symptomatic breast cancers are downloaded from the UK cancer registries. The data are then sent to individual surgeons for validation. Validated data are returned to the WMCIU for analysis. In this report, where altered data were returned by surgeons in England, Wales and Northern Ireland, these have been used in the analysis in preference to the original cancer registration data. Data for Scotland were provided by the Information and Statistics Division Scotland which has managed the Scottish Cancer Registration scheme since 1997.

NHSBSP/ABS validated data – Data for the UK NHS Breast Screening Programme (NHSBSP) and Association of Breast Surgery (ABS) audit of screen-detected breast cancer are initially downloaded from the National Breast Screening System (NBSS) or other breast screening computer systems. Data are checked and signed off by the responsible surgeons and the regional breast screening QA reference centres prior to their inclusion in the audit. These data were used to assign a screen-detected flag to cases.

Cohort Characteristics

Key Findings

- In the UK in 2007, 50,286 new cases of invasive and non/micro-invasive breast cancer were registered. 44,782 cases were invasive (89%) and 5,490 cases (11%) were non-invasive or micro-invasive.
- 292 new breast cancers (0.6%) were diagnosed in men.
- 81% of new breast cancers were diagnosed in patients aged 50 and over.
- In women, 32% of all breast cancers and 56% of breast cancers diagnosed between the ages of 50 and 69 years were screen-detected.
- There was a marked relationship between deprivation and breast cancer incidence; with only 14-17% of breast cancers being diagnosed in people in the most deprived quintile compared with 20-23% in the least deprived quintile.
- Deprivation profiles varied to some extent with age at diagnosis, with patients aged less than 40 being more evenly distributed across the deprivation quintiles than other age groups, and significantly fewer patients aged 75 and over in the least deprived quintile (19% compared to 24% under 75).
- In women known to be Black, 46% of breast cancers were diagnosed under the age of 50. Conversely, 30% of breast cancers in women known to be White were diagnosed in those aged 70 and over.
- 75% of women with breast cancer known to be Black and 62% of women known to be Asian were in the two most deprived quintiles (Q1 and Q2). In contrast, 45% of women with breast cancer known to be White and 40% of women known to be Chinese were in the two least deprived quintiles (Q4 and Q5).
- Only 20% of women known to be Black had screen-detected breast cancers compared to 32% of all women.

Country Profile

A total of 50,286 cases of invasive and non/micro-invasive breast cancer diagnosed in the UK in 2007 are included in this report. Of these, 83.4% were diagnosed in England, 2.5% in Northern Ireland, 8.8% in Scotland and 5.3% in

Wales. 49,994 breast cancers were diagnosed in women (99.4%) and 292 in men (0.6%). Details of the number of breast cancers diagnosed in women and men in each English region and Celtic country are given in Table 1.

| English Region/ Celtic Country | Population Covered (million) | No. Breast Cancers in 2007 | | | % Breast Cancers in UK |
|-----------------------------------|---------------------------------|----------------------------|---------------|---------------|---------------------------|
| | | Men | Women | Total | |
| Eastern | 5.66 | 37 | 4,800 | 4,837 | 9.6% |
| North West | 6.58 | 31 | 5,394 | 5,425 | 10.8% |
| Northern & Yorkshire | 6.76 | 34 | 5,492 | 5,526 | 11.0% |
| Oxford | 2.86 | 10 | 2,236 | 2,246 | 4.5% |
| South West | 7.02 | 43 | 6,698 | 6,741 | 13.4% |
| Thames | 11.84 | 47 | 8,434 | 8,481 | 16.9% |
| Trent | 4.99 | 25 | 3,945 | 3,970 | 7.9% |
| West Midlands | 5.38 | 25 | 4,710 | 4,735 | 9.4% |
| England | 51.09 | 252 | 41,709 | 41,961 | 83.4% |
| Northern Ireland | 1.76 | 3 | 1,269 | 1,272 | 2.5% |
| Scotland | 5.14 | 26 | 4,382 | 4,408 | 8.8% |
| Wales | 2.98 | 11 | 2,634 | 2,645 | 5.3% |
| UK | 60.98 | 292 | 49,994 | 50,286 | |

Table 1: English region and Celtic country profiles

Cohort Characteristics




Figure 1: Proportion of breast cancer cases in each region in each deprivation group

In England as a whole, there was a marked relationship between deprivation and breast cancer incidence; with only 14% of breast cancers being diagnosed in people in the most deprived quintile compared with 23% in the least deprived quintile (Figure 1). Each deprivation quintile covers 20% of the English population. The proportion of breast cancers diagnosed in the most deprived group is significantly smaller than the expected 20% and the proportion in the least deprived group significantly in excess. This implies that people in the least deprived population are more likely to be diagnosed

with breast cancer. The large regional differences in the proportions of breast cancer diagnosed in each deprivation quintile within England reflect the marked variation in deprivation levels between the regions; the Eastern, Oxford and South West regions being less deprived compared to the rest of the England. The proportions of breast cancer in the five deprivation groups in the Celtic countries are similar to those in England; with a higher proportion diagnosed in people in the least deprived quintile (20%-22% compared with 16-17% in the most deprived quintile).

Age, Gender and Deprivation

The cohort of patients diagnosed with breast cancer in 2007 had an age distribution ranging from 17 to 103 years. 81% of breast cancers were diagnosed in patients aged 50 and over (Table 2). The median age at diagnosis in men was higher than in women (72 compared to 62); 54% of breast cancers were diagnosed in men aged 70 and over compared with 31% in women. Men diagnosed with breast cancer were less likely to have non-invasive

or micro-invasive tumours (7.9% compared to 10.9%), but this difference was not statistically significant.

Because of the small number of micro-invasive breast cancers and because they are treated in a similar way, in the remainder of this report, non-invasive and micro-invasive breast cancers have been combined into a single non/micro-invasive category.

Cohort Characteristics

| Characteristic | Women | | Men | | Total | |
|---------------------------------|--------|-----|-----|-----|--------|------|
| | No. | % | No. | % | No. | % |
| All breast cancers | 49,994 | 99% | 292 | 1% | 50,286 | 100% |
| Age at Diagnosis (years) | | | | | | |
| < 30 | 193 | 0% | 3 | 1% | 196 | 0% |
| 30 - 49 | 9,136 | 18% | 18 | 6% | 9,154 | 18% |
| 50 - 69 | 25,124 | 50% | 113 | 39% | 25,237 | 50% |
| 70 - 89 | 14,154 | 28% | 149 | 51% | 14,303 | 28% |
| 90 + | 1,387 | 3% | 9 | 3% | 1,396 | 3% |
| Deprivation Quintile | | | | | | |
| Q1 Most deprived | 7,349 | 15% | 37 | 13% | 7,386 | 15% |
| Q2 | 9,202 | 18% | 67 | 23% | 9,269 | 18% |
| Q3 | 10,590 | 21% | 54 | 18% | 10,644 | 21% |
| Q4 | 11,393 | 23% | 75 | 26% | 11,468 | 23% |
| Q5 Least deprived | 11,262 | 22% | 59 | 20% | 11,321 | 23% |
| Unknown | 198 | 0% | | 0% | 198 | 0% |
| Tumour Invasive Status | | | | | | |
| Invasive | 44,513 | 89% | 269 | 92% | 44,782 | 89% |
| Micro-invasive | 258 | 1% | 2 | 0% | 260 | 1% |
| Non-invasive | 5,209 | 10% | 21 | 7% | 5,230 | 10% |
| Unknown | 14 | 0% | | 0% | 14 | 0% |

Table 2: Characteristics of breast cancers diagnosed in women and men

The deprivation profiles for breast cancers diagnosed in men and women were similar, with 46% and 45% respectively being diagnosed in patients in the two least deprived quintiles (Q4 and Q5) (Table 2). Deprivation profiles varied to some extent with age at diagnosis, with patients aged less than 40 being more evenly distributed across the deprivation quintiles than other age groups, and significantly fewer patients aged 75 and over in the least deprived quintile (19% compared to 24% under 75).




Figure 2: Variation in deprivation status with age at diagnosis for women diagnosed with breast cancer in 2007

Cohort Characteristics

The flat deprivation profile for women aged less than 40 seen in Figure 2 may reflect the relatively high proportion of these cancers which are familial rather than being linked to the lifestyle factors which are believed to influence the incidence of sporadic breast cancer. The decrease in the proportion of breast cancers in the two least deprived quintiles in women aged 75 and over may be due to the earlier diagnosis by the NHS

Route of Presentation

Of the 49,994 female breast cancers diagnosed in 2007, 15,783 (32%) were detected by the NHS Breast Screening Programme. The majority were diagnosed in women in the screening age range (50-70 years in 2007), with a small number detected in women aged less than 50 who were called early to screening to ensure that they would be invited before they were aged 53, and 416 in women aged 75 years and over who had self-referred for screening.

The age distribution of screen-detected and symptomatic breast cancers diagnosed in women in the UK in 2007 is shown in Figure 3. 50% of all breast cancers were

Breast Screening Programme of breast cancers that would have occurred in this age group.

Because of the relatively small number of breast cancers diagnosed in men, the remainder of this report includes only breast cancers diagnosed in women. A future publication on breast cancer in men is planned which will include data aggregated over a number of years.

diagnosed in women between the ages of 50 and 69 years, and 56% of these were screen-detected. Although 31% of all breast cancers in women were diagnosed in patients aged 70 and over, only 7% of these were screen-detected. 2% of breast cancers diagnosed in women aged less than 50 were detected via screening. In England, 5% of female breast cancer patients (2,176 women) were 47-49 years old at diagnosis, and a further 5% (1,914 women) were 71-73 years old. These groups of women are covered by the extension of the NHS Breast Screening Programme outlined in the Cancer Reform Strategy and therefore, in future, more breast cancers in these age groups may be screen-detected.




Figure 3: Age profile and route of presentation for women diagnosed with breast cancer in 2007

Cohort Characteristics

The proportion of screen-detected breast cancers was significantly lower in the Thames region than in the rest of the UK (27% compared to 32%). This probably partly reflects the relatively high proportion of women in minority ethnic groups in this region (15% in Thames and 3% in the rest of England) as these women are known to be less likely to accept their invitation to attend for screening⁹. The proportion of screen-detected

cancers in Northern Ireland was also significantly lower than the rest of the UK (25% compared to 32%). This is partly because in 2007 the Northern Ireland Breast Screening Programme had not been expanded to cover women aged 65-70 years, with the result that only 49% of cancers diagnosed in this age group were screen-detected compared with 56% in the rest of the UK where the expansion had occurred.

Ethnicity

Ethnicity data were not available for the Celtic countries. 33,250 (79.7%) of the women diagnosed with breast cancer in England had a recorded ethnic group. Of the 8,459 cases where ethnicity was unknown, half were

women aged 65 years and over. This is in part because ethnicity is obtained from Hospital Episode Statistics data and older women are less likely to have surgery and thus a hospital admission record.

| Characteristic | Ethnic Group | | | | | | | Total |
|---------------------------------|--------------|-------|-------|---------|-------|-------|---------|--------|
| | White | Asian | Black | Chinese | Mixed | Other | Unknown | |
| Number of cases | 31,443 | 820 | 505 | 85 | 114 | 283 | 8,459 | 41,709 |
| % cases | 75% | 2% | 1.2% | 0.2% | 0.3% | 0.7% | 20% | |
| Age at Diagnosis (years) | | | | | | | | |
| < 30 | 0% | 1% | 1% | 1% | 1% | 0% | 0% | 0% |
| 30 - 49 | 18% | 27% | 45% | 31% | 31% | 34% | 16% | 18% |
| 50 - 69 | 52% | 59% | 39% | 59% | 52% | 48% | 44% | 50% |
| 70 - 89 | 28% | 12% | 14% | 8% | 16% | 17% | 34% | 28% |
| 90 + | 2% | 0% | 1% | 1% | 1% | 0% | 6% | 3% |
| Deprivation Quintile | | | | | | | | |
| Q1 Most deprived | 14% | 41% | 50% | 15% | 22% | 30% | 11% | 14% |
| Q2 | 19% | 21% | 25% | 25% | 28% | 23% | 17% | 18% |
| Q3 | 22% | 16% | 12% | 20% | 18% | 17% | 21% | 21% |
| Q4 | 23% | 9% | 6% | 28% | 18% | 17% | 24% | 23% |
| Q5 Least deprived | 22% | 13% | 7% | 12% | 13% | 12% | 27% | 23% |
| Unknown | 0% | 0% | 0% | 0% | 1% | 0% | 1% | 0% |
| Tumour Invasive Status | | | | | | | | |
| Invasive | 89% | 89% | 87% | 81% | 87% | 89% | 88% | 89% |
| Non/micro-invasive | 11% | 11% | 13% | 19% | 13% | 11% | 12% | 11% |
| Unknown | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% |
| Presentation Route | | | | | | | | |
| Screen-detected | 33% | 31% | 20% | 35% | 32% | 28% | 26% | 32% |
| Symptomatic | 67% | 69% | 80% | 65% | 68% | 72% | 74% | 68% |

Table 3: Variations in age, deprivation status, invasive status and presentation route with ethnic group for women diagnosed with breast cancer in England in 2007

Cohort Characteristics

In women of known Black ethnicity, 46% of breast cancers were diagnosed under the age of 50, compared to 32% in women known to be Chinese or of Mixed ethnicity, 28% in women known to be Asian and 18% in women known to be White. Conversely, 30% of breast cancers in women known to be White were diagnosed in those aged 70 and over compared with 9-17% of women known to be Chinese, Asian, Black or of Mixed ethnicity. These differences will in part reflect differences in age distribution in the minority ethnic groups compared to the White population and have not been examined further in this report.

75% of women diagnosed with breast cancer known to be Black and 62% of women known to be Asian were in the two most deprived quintiles (Q1 and Q2) compared to 33% of women known to be White, 40% of women known to be Chinese and 50% of women known to be of Mixed ethnicity. In contrast, 45% of women diagnosed with breast cancer known to be White and 40% of women known to be Chinese were in the two least deprived quintiles (Q4 and Q5) compared with 32% of women known to be of Mixed ethnicity, 22% of women known to be Asian and 13% of women known to be

Black. As noted in the first *All Breast Cancer Report*⁵, women known to be Chinese had a higher proportion of non/micro-invasive cancers (19% compared to 11% in the rest of England), but this difference is not statistically significant.

Only 20% of women known to be Black had screen-detected breast cancers compared to 32% of all women. The low proportion of women of known Black ethnicity with screen-detected breast cancer may in part be explained by the relatively high proportion of cancers (46%) diagnosed in these women under the age of 50. However, there were also relatively high proportions of breast cancers diagnosed in women under the age of 50 known to be Asian (28%), Chinese (32%) or of Mixed ethnicity (32%), and 31-35% of their cancers were screen-detected so this cannot be the only reason.

The remainder of this report does not include analyses for individual ethnic groups. A future publication on breast cancer and ethnicity is planned which will include data aggregated over a number of years to allow more meaningful analysis of the differences between the minority ethnic groups.

Tumour Characteristics

Key Findings

- Of the surgically treated women diagnosed with invasive breast cancer with known tumour characteristics, 58% had small cancers with an invasive tumour size of 20mm or less, 17% were diagnosed with grade 1 tumours, 62% had lymph node negative tumours, 39% had tumours in the excellent (EPG) and good (GPG) Nottingham Prognostic Index groups, 29% had vascular invasion, 84% had ER positive tumours and 15% had HER2 positive tumours.
- Screen-detected invasive breast cancers were more likely to be smaller in size, of a lower grade, node negative and in the EPG and GPG prognostic groups. These cancers were also less likely to have vascular invasion, more likely to be ER positive and less likely to be HER2 positive. These differences between symptomatic and screen-detected cancers were evident across all ages and deprivation quintiles.
- Older women diagnosed symptomatically, were more likely to have node negative cancers and EPG/GPG cancers, and less likely to have small cancers, HER2 positive cancers and vascular invasion present. Older women diagnosed via screening were also more likely to have node negative cancers, and less likely to have HER2 positive cancers and vascular invasion.
- Women aged less than 40 diagnosed symptomatically were less likely to have grade 1 tumours and cancers in the EPG and GPG prognostic groups, and more likely to have vascular invasion present.
- For symptomatic invasive breast cancers, the proportions of small cancers, node negative cancers, cancers in the EPG and GPG prognostic groups and ER positive cancers decreased with increasing deprivation. Equivalent trends were not apparent for screen-detected invasive breast cancers indicating that screening reduces the observable inequalities associated with deprivation.

Tumour Characteristics

Tumour characteristics data were available for the whole of the UK. Tables summarising the completeness of each data item for invasive breast cancers diagnosed in each English region and Celtic country are provided in Appendix 3. Variations in tumour size, degree of spread, aggressiveness and receptor status are considered in

terms of age at diagnosis, deprivation and presentation route in women diagnosed with invasive breast cancer in 2007. Breast cancers in men have been excluded from this section and the remainder of the report, as has the effect of ethnicity on tumour characteristics and treatment outcomes.

Invasive Tumour Size

In the UK, invasive tumour size was known for 76% of all invasive cancers and for 89% of surgically treated invasive cancers. The former varied from 41% in the

Trent region to 90% in the Eastern region, and the latter from 53% in the Trent region to 98% in the West Midlands region (Appendix 3).

Tumour Characteristics

| Age Band (years) | Invasive Cancers Tumour Size 20mm or less | | | |
|------------------|---|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | Total no. cases |
| <40 | - | 48.4% | 48.4% | 1462 |
| 40-49 | 74.4% | 49.2% | 49.8% | 4864 |
| 50-59 | 77.4% | 47.8% | 64.3% | 8480 |
| 60-69 | 79.2% | 49.4% | 68.1% | 9700 |
| 70-79 | 79.4% | 41.1% | 49.7% | 5392 |
| 80+ | 69.6% | 36.2% | 37.3% | 2216 |
| All ages | 78.4% | 45.8% | 58.2% | 32114 |

| Deprivation Quintile | Invasive Cancers Tumour Size 20mm or less | | | |
|----------------------|---|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | Total no. cases |
| Q1 Most deprived | 76.7% | 42.9% | 54.7% | 4589 |
| Q2 | 78.9% | 44.1% | 56.6% | 5826 |
| Q3 | 79.2% | 45.1% | 57.9% | 6760 |
| Q4 | 78.7% | 48.0% | 60.2% | 7388 |
| Q5 Least deprived | 78.2% | 47.5% | 60.0% | 7478 |
| All cases | 78.4% | 45.8% | 58.2% | 32041 |

Tables 4a and 4b: Variation in the proportion of surgically treated small (maximum diameter 20mm or less) invasive breast cancers with a) age and route of presentation and b) deprivation quintile and route of presentation expressed as a proportion of the cancers with known invasive tumour size

Overall, 58% of surgically treated women diagnosed with invasive breast cancer in 2007 had small cancers with an invasive tumour size 20mm or less (Table 4a). The proportion of screen-detected cancers with an invasive tumour size 20mm or less was significantly higher than that for symptomatic cancers (78% compared to 46%). Women aged 70 years and older who presented symptomatically were less likely to be diagnosed with a small cancer (41% in women aged 70-79 and 36% in women aged 80 or over). Women aged 80 and over were less likely to have small tumours regardless of their route of presentation. This difference was statistically significant for symptomatic but not screen-detected cases.

Invasive Tumour Grade

Invasive tumour grade was known for 90% of all invasive cancers and for 97% of surgically treated invasive cancers. The former varied from 85% in Northern Ireland

The proportions of surgically treated women with small (20mm or less) invasive breast cancers were lowest in the most deprived quintile for both screen-detected and symptomatic cancers (Table 4b), but in every quintile the proportion of small (20mm or less) breast cancers was significantly higher for screen-detected cancers. For symptomatic cancers there is a statistically significant relationship between the proportion of small (20mm or less) invasive cancers and deprivation. The proportion of small (20mm or less) invasive cancers increased from 43% in the most deprived quintile to 48% in the two least deprived quintiles. This relationship is not apparent for screen-detected cancers.

to 94% in the Oxford region, and the latter from 93% in Northern Ireland to 99% in the Oxford and West Midlands regions (Appendix 3).

Tumour Characteristics

| Age Band (years) | Invasive Cancers Grade 1 | | | Total no. cases |
|------------------|--------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 5.9% | 5.9% | 1713 |
| 40-49 | 28.6% | 10.7% | 11.1% | 5613 |
| 50-59 | 28.5% | 11.3% | 20.4% | 8938 |
| 60-69 | 26.3% | 11.0% | 20.1% | 10129 |
| 70-79 | 26.6% | 11.5% | 14.6% | 5836 |
| 80+ | 21.7% | 11.5% | 11.8% | 2510 |
| All ages | 27.2% | 10.7% | 16.5% | 34739 |

| Deprivation Quintile | Invasive Cancers Grade 1 | | | Total no. cases |
|----------------------|--------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 24.8% | 10.6% | 15.2% | 4957 |
| Q2 | 26.1% | 10.7% | 15.7% | 6341 |
| Q3 | 29.2% | 10.4% | 17.0% | 7275 |
| Q4 | 27.0% | 10.7% | 16.6% | 7991 |
| Q5 Least deprived | 27.7% | 11.2% | 17.4% | 8100 |
| All cases | 27.2% | 10.7% | 16.5% | 34664 |

Tables 5a and 5b: Variation in the proportion of surgically treated grade 1 invasive breast cancers with a) age and route of presentation and b) deprivation quintile and route of presentation expressed as a proportion of the cancers with known grade

Grade 1 cancers are known to have a better prognosis than higher grade disease. Overall, 17% of surgically treated women diagnosed with invasive breast cancer in 2007 had a grade 1 cancer (Table 5a). The proportion of screen-detected grade 1 cancers was significantly higher than that for symptomatic cancers (27% compared to 11%). The proportion of grade 1 cancers varied with age at diagnosis for both presentation routes. Women aged less than 40 with symptomatic cancers had a

significantly smaller proportion of grade 1 tumours (6%), as did women aged 80 or over with screen-detected cancers (22%) but, for the latter age group, this variation was not statistically significant. The proportions of surgically treated women with grade 1 cancers were significantly higher for screen-detected tumours in all deprivation quintiles (Table 5b). There was no overall trend across the deprivation quintiles for either screen-detected or symptomatic cancers.

Nodal Status

Within the UK, the nodal status of invasive cancers was known for 66% of all invasive cancers and for 80% of surgically treated invasive cancers. The former varied

from 45% in Wales to 84% in Northern Ireland, and the latter from 52% in the Trent region to 96% in the West Midlands region and Scotland (Appendix 3).

Tumour Characteristics

| Age Band (years) | Invasive Cancers Negative Nodal Status | | | Total no. cases |
|------------------|--|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 45.8% | 45.8% | 1285 |
| 40-49 | 74.6% | 47.8% | 48.6% | 4235 |
| 50-59 | 75.1% | 48.9% | 64.6% | 7838 |
| 60-69 | 79.0% | 52.0% | 70.0% | 9037 |
| 70-79 | 79.8% | 52.9% | 59.8% | 4621 |
| 80+ | 71.0% | 50.9% | 51.6% | 1613 |
| All ages | 77.5% | 50.0% | 61.6% | 28629 |

| Deprivation Quintile | Invasive Cancers Negative Nodal Status | | | Total no. cases |
|----------------------|--|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 75.4% | 47.6% | 58.7% | 3959 |
| Q2 | 77.4% | 48.4% | 60.1% | 5100 |
| Q3 | 79.8% | 50.1% | 62.3% | 6096 |
| Q4 | 76.6% | 51.3% | 62.4% | 6648 |
| Q5 Least deprived | 77.6% | 50.9% | 62.8% | 6758 |
| All cases | 77.5% | 49.9% | 61.6% | 28561 |

Tables 6a and 6b: Variation in the proportion of surgically treated node negative invasive breast cancers with a) age and route of presentation and b) deprivation quintile and route of presentation expressed as a proportion of the cancers with known nodal status

Node negative breast cancers are known to have a better prognosis than node positive cancers. In 2007, 62% of surgically treated women diagnosed with invasive breast cancer had node negative cancers (Table 6a). The proportion of screen-detected node negative cancers was significantly higher than the proportion of symptomatic cancers (78% compared to 50%). For symptomatic cancers there is a statistically significant increase in the proportion of node negative cancers with increasing age at diagnosis. Node negative cancers increased from 46% in women aged less than 40 to 53% in those aged 70-79. A similar statistically significant relationship is also apparent for screen-detected cancers up to the age of 70-79 years. Women aged 80 and over

with screen-detected cancers had a smaller proportion of node negative cancers (71%), but this difference is not statistically significant.

The proportions of surgically treated women with node negative invasive breast cancers were significantly higher for screen-detected cancers in all deprivation quintiles (Table 6b). For surgically treated women with symptomatic cancers, there is a statistically significant increase in the proportion of node negative cancers with decreasing deprivation. The proportion of node negative cancers increased from 48% in the most deprived quintile to 51% in the two least deprived quintiles. This relationship is not apparent for screen-detected cancers.

Nottingham Prognostic Index (NPI)

The Nottingham Prognostic Index (NPI)¹⁰ is used to determine the prognosis of surgically treated invasive breast cancers. An NPI score is calculated using three pathological criteria: invasive size, number of involved nodes and tumour grade. The scores can be grouped into five distinct prognostic groups: Excellent (EPG), Good (GPG), Moderate 1 (MPG1), Moderate 2 (MPG2)

and Poor (PPG). In the UK as a whole, the Nottingham Prognostic Index was known for 63% of all invasive cancers and 76% of surgically treated invasive cancers. The former varied from 40% in Trent to 79% in the West Midlands region, and the latter from 51% in the Trent region to 95% in the West Midlands region (Appendix 3).

Tumour Characteristics

| Age Band (years) | Invasive Cancers EPG/GPG | | | Total no. cases |
|------------------|--------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 14.8% | 14.8% | 1149 |
| 40-49 | 60.0% | 23.6% | 24.8% | 3907 |
| 50-59 | 56.5% | 23.4% | 43.5% | 7579 |
| 60-69 | 60.2% | 25.6% | 49.0% | 8792 |
| 70-79 | 62.2% | 26.0% | 35.4% | 4476 |
| 80+ | 41.9% | 25.5% | 26.2% | 1562 |
| All ages | 58.9% | 24.0% | 39.1% | 27465 |

| Deprivation Quintile | Invasive Cancers EPG/GPG | | | Total no. cases |
|----------------------|--------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 54.6% | 20.1% | 34.3% | 3808 |
| Q2 | 58.7% | 22.5% | 37.5% | 4889 |
| Q3 | 61.6% | 24.4% | 40.1% | 5872 |
| Q4 | 57.8% | 25.9% | 40.2% | 6370 |
| Q5 Least deprived | 60.0% | 25.1% | 41.1% | 6461 |
| All cases | 58.9% | 23.9% | 39.1% | 27400 |

Tables 7a and 7b: Variation in the proportion of surgically treated invasive breast cancers in the Excellent (EPG) and Good (GPG) NPI groups with a) age and route of presentation and b) deprivation quintile and route of presentation expressed as a proportion of the cancers with known NPI

Overall, 39% of surgically treated women diagnosed with invasive breast cancer in 2007 had EPG or GPG cancers (Table 7a). The proportion of EPG/GPG cancers was significantly higher in the screen-detected cohort than for symptomatic cancers (59% compared to 24%). For symptomatic cancers, there is a statistically significant relationship between the proportion of EPG/GPG cancers and age at diagnosis; older women having a higher proportion of EPG/GPG cancers. Women aged less than 40 had the lowest proportion of EPG/GPG cancers (15%). Women aged 80 or over with screen-detected cancers had a significantly smaller proportion of EPG/GPG cancers than women with screen-detected cancers in the other age groups, but this was higher than in women in the same age group with symptomatic cancers.

Vascular Invasion

Vascular invasion data were not available for Northern Ireland and Scotland. In England and Wales, vascular invasion was known for 37% of all invasive cancers and for 44% of surgically treated invasive cancers. The

former varied from 9% in the Thames region to 84% in the Oxford region, and the latter from 10% in the Thames region to 93% in the West Midlands region (Appendix 3).

Across all deprivation quintiles, the proportions of surgically treated women with EPG/GPG cancers were significantly higher for screen-detected breast cancers (Table 7b). For women with symptomatic cancers, the proportion of EPG/GPG cancers increased from 20% in the most deprived quintile to 25% in the least deprived quintile, and women in the most deprived quintile were significantly less likely to have EPG/GPG cancers. Although the overall relationship between the proportion of early stage cancers and deprivation was not apparent for screen-detected cancers, women in the most deprived quintile with screen-detected cancers did have a significantly lower proportion of EPG/GPG cancers.

Tumour Characteristics

| Age Band (years) | Invasive Cancers Vascular Invasion Present | | | Total no. cases |
|------------------|---|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 40.8% | 40.8% | 732 |
| 40-49 | 17.0% | 35.0% | 34.7% | 2442 |
| 50-59 | 16.8% | 33.1% | 25.7% | 3294 |
| 60-69 | 16.3% | 32.0% | 24.0% | 3781 |
| 70-79 | 15.6% | 32.0% | 29.4% | 2562 |
| 80+ | 5.0% | 30.4% | 30.0% | 1137 |
| All ages | 16.4% | 33.4% | 28.6% | 13948 |

| Deprivation Quintile | Invasive Cancers Vascular Invasion Present | | | Total no. cases |
|----------------------|---|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 17.9% | 32.9% | 29.5% | 1709 |
| Q2 | 16.1% | 34.3% | 29.6% | 2379 |
| Q3 | 15.0% | 33.0% | 27.9% | 3073 |
| Q4 | 15.7% | 32.1% | 27.5% | 3347 |
| Q5 Least deprived | 17.7% | 35.0% | 29.6% | 3386 |
| All cases | 16.4% | 33.5% | 28.7% | 13894 |

Tables 8a and 8b: Variation in the proportion of surgically treated invasive breast cancers with vascular invasion present with a) age and route of presentation and b) deprivation quintile and route of presentation expressed as a proportion of cancers with known vascular invasion

Overall, 29% of surgically treated women diagnosed with invasive breast cancer in 2007 had vascular invasion present. The proportion of screen-detected cancers with vascular invasion present was significantly lower than for that for symptomatic cancers (16% compared to 33%) (Table 8a). The proportions of cancers with vascular invasion present decreased with increasing age for both presentation routes. Women aged less than 40 with symptomatic cancer had a significantly higher proportion with vascular invasion present (41% compared with 29% overall). Women aged 80 and over,

with screen-detected or symptomatic cancer, had a lower proportion of cancers with vascular invasion present, but these differences are not statistically significant.

The proportions of surgically treated women with invasive breast cancers with vascular invasion present were significantly lower for screen-detected cancers in all deprivation quintiles, but there was no clear trend in the presence of vascular invasion across the deprivation quintiles for either presentation route (Table 8b).

Oestrogen Receptor Status (ER Status)

Data on ER status were not available for Northern Ireland. For the remaining countries, ER status was known for 56% of all invasive cancers and 61% of surgically treated invasive cancers. The former varied

from 32% in the Thames region to 96% in the West Midlands region, and the latter from 37% in the Thames region to 99% in the West Midlands region (Appendix 3).

Tumour Characteristics

| Age Band (years) | Invasive Cancers ER Positive | | | Total no. cases |
|------------------|------------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 67.9% | 67.9% | 724 |
| 40-49 | 95.0% | 79.2% | 80.0% | 2484 |
| 50-59 | 89.3% | 73.1% | 84.4% | 6257 |
| 60-69 | 90.2% | 76.4% | 86.9% | 7593 |
| 70-79 | 91.4% | 79.7% | 83.8% | 3233 |
| 80+ | 91.0% | 78.4% | 79.1% | 1162 |
| All ages | 90.0% | 76.7% | 83.8% | 21453 |

| Deprivation Quintile | Invasive Cancers ER Positive | | | Total no. cases |
|----------------------|------------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 88.5% | 75.5% | 82.1% | 3053 |
| Q2 | 89.2% | 74.9% | 82.3% | 3833 |
| Q3 | 90.8% | 76.9% | 84.1% | 4563 |
| Q4 | 90.8% | 76.7% | 84.5% | 4971 |
| Q5 Least deprived | 89.9% | 78.9% | 85.2% | 4970 |
| All cases | 90.0% | 76.7% | 83.8% | 21390 |

Tables 9a and 9b: Variation in the proportion of surgically treated ER positive invasive breast cancers with a) age and route of presentation and b) deprivation quintile and route of presentation expressed as a proportion of the cancers with known ER status

Overall, 84% of surgically treated women diagnosed with invasive breast cancer in 2007 had ER positive tumours (Table 9a). The proportion of screen-detected ER positive cancers was significantly higher than that for symptomatic cancers overall (90% compared to 77%) and in every age group. There is a non linear relationship between age and ER positive status for both screen-detected and symptomatic cancers. For screen-detected cases the highest proportion of ER positive cancers was in the relatively small cohort of women aged 40-49, but this difference is not statistically significant.

Human Epidermal Growth Factor Receptor 2 Status (HER2 Status)

Data on HER2 status were not available for Northern Ireland and Scotland. In England and Wales, HER2 status was known for only 43% of all invasive cancers and for only 50% of surgically treated invasive cancers.

The proportions of surgically treated women with ER positive invasive breast cancers were significantly higher for screen-detected cancers in all deprivation quintiles (Table 9b). For symptomatic cancers, the proportion of ER positive cancers increased significantly with decreasing deprivation. The proportion of ER positive invasive breast cancers was 75% in the most deprived quintile and 79% in the least deprived quintile. This relationship is not apparent for screen-detected cancers.

The former varied from 26% in the Thames region to 87% in the West Midlands region, and the latter from 30% in the Thames region to 92% in the West Midlands region (Appendix 3).

Tumour Characteristics

| Age Band (years) | Invasive Cancers HER2 Positive | | | Total no. cases |
|------------------|--------------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 26.1% | 26.1% | 532 |
| 40-49 | 20.5% | 19.0% | 19.1% | 1789 |
| 50-59 | 14.9% | 19.0% | 16.1% | 4702 |
| 60-69 | 11.8% | 18.3% | 13.2% | 5771 |
| 70-79 | 8.8% | 14.9% | 12.6% | 2257 |
| 80+ | 5.0% | 12.8% | 12.4% | 764 |
| All ages | 12.8% | 17.9% | 15.0% | 15815 |

| Deprivation Quintile | Invasive Cancers HER2 Positive | | | Total no. cases |
|----------------------|--------------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 11.2% | 18.1% | 6.0% | 2276 |
| Q2 | 13.3% | 17.2% | 7.3% | 2821 |
| Q3 | 12.0% | 16.5% | 6.6% | 3282 |
| Q4 | 13.7% | 19.6% | 7.9% | 3690 |
| Q5 Least deprived | 13.0% | 18.3% | 7.7% | 3694 |
| All cases | 12.8% | 18.0% | 7.2% | 15763 |

Tables 10a and 10b: Variation in the proportion of surgically treated HER2 positive invasive breast cancers with a) age and route of presentation and b) deprivation quintile and route of presentation expressed as a proportion of the cancers with known HER2 status

Overall, 15% of surgically treated women diagnosed with invasive breast cancer in 2007 had HER2 positive tumours (Table 10a). This is significantly lower than the early data that indicated over-expression levels of 20-25%¹¹, but it is in line with more recent publications that have revised the figure to as low as 9% in some series^{12,13}. Overall, the proportion of screen-detected HER2 positive invasive breast cancers was significantly lower than for that for symptomatic cancers (13% compared to 18%). There are significant inverse relationships between increasing

age and HER2 positivity for screen-detected and symptomatic cancers; invasive breast cancers in older patients being significantly less likely to be HER2 positive.

The proportions of surgically treated women with HER2 positive invasive breast cancers were significantly lower for screen-detected cancers compared to symptomatic cancers in all deprivation quintiles (Table 10b). However, there was no significant variation between deprivation quintiles in the proportions of HER2 positivity for screen-detected or symptomatic cancers.

Surgical Treatment Outcomes

Key Findings

- 82% of women diagnosed with breast cancer in the UK in 2007 had surgery. Significantly more women with screen-detected breast cancer had surgery recorded (98% compared to 75% of women presenting symptomatically).
- For symptomatic breast cancers, surgical treatment decreased with age at diagnosis with only 74% of women aged 70-79 and 39% of women aged 80 and above having surgical treatment compared with 90% of women aged under 50.
- Women aged 50-69 in the most deprived group with symptomatic breast cancers were less likely to have surgery than women in the least deprived group (72% compared to 78%). This effect was not seen in women with screen-detected breast cancer.
- Overall 57% all women with surgically treated breast cancer had a breast conserving procedure. The mastectomy rate was lower for women with screen-detected breast cancer (27% compared to 53%), and a higher proportion of women with screen-detected breast cancer had breast conserving surgery in every age band.
- For women with symptomatic but not screen-detected cancer, the likelihood of having breast conserving surgery decreased with increasing deprivation.
- Although the overall mastectomy rate for invasive breast cancers was significantly higher than that for non/micro-invasive breast cancers (44% compared to 38%), for screen-detected cases, the mastectomy rate for non/micro-invasive breast cancers was significantly higher (30% compared to 26%). The latter may reflect the higher rate of immediate reconstruction for non/micro-invasive breast cancers which form a much higher proportion of screen-detected cases (20% compared to 8% of symptomatic cancers).
- In the UK as a whole for all breast cancers, the MX:BCS ratio was 1:1.3. This is indicative of a higher level of breast conserving surgery (BCS) than anticipated by the national reference figure of 1:1. The MX:BCS ratio was lower (i.e. the proportion of cases treated with breast conserving surgery was higher) for screen-detected breast cancers than for symptomatic breast cancers and for non/micro-invasive breast cancers than for invasive breast cancers.
- Overall, 11% of female mastectomy patients had an immediate reconstruction. A greater proportion of non/micro-invasive cases had a reconstruction (27% versus 10% of invasive cases). Women with screen-detected cancer were more likely to have reconstruction (16% versus 10%), and for these patients, reconstruction was significantly more likely to be undertaken in those from a less deprived background.
- Overall, 32% of women had a Sentinel Lymph Node Biopsy (SLNB). Women with poor prognostic group cancers and older women were less likely to have this procedure.
- The median length of stay in hospital for women having a mastectomy was 4 days compared to 2 days for those having breast conserving surgery. Screening patients had an overall shorter length of stay mainly because more were treated with breast conserving surgery. Older women and more deprived women had a longer length of stay.
- Repeat operation rates were higher for women with non/micro-invasive cancer (25% compared with 14% for invasive cancers), for screen-detected cancers (18%) and for younger women (19%).
- 860 women (2%) with breast cancer died within 30 days of their diagnosis. 51% of these were death certificate only (DCO) registrations. Of the non-DCO cases who died within 30 days, 64% died of breast cancer. A significantly higher proportion of women in the most deprived quintile died within 30 days of their diagnosis of breast cancer.
- In 2007 in England, Wales and Northern Ireland, 524 surgeons operated on female breast cancer patients. 129 (27%) surgeons had a caseload of less than 30 patients. 27 surgeons had a significantly high mastectomy rate and 44 surgeons a significantly low mastectomy rate.

Surgical Treatment Outcomes

Surgical Treatment

Overall, 82% of the women diagnosed with breast cancer in the UK in 2007 had surgery recorded. This varied from 76% in Northern Ireland to 88% in the Oxford region. Significantly more women with screen-detected breast cancer had surgery recorded (98% compared to 75% of women presenting symptomatically). This difference is likely to be due in part to the better prognosis of screen-detected cancers and to the younger age of the majority of the women with screen-detected cancer. Also, as symptomatic women were more likely to be deprived, they may have had more co-morbid conditions which could have affected their eligibility for surgery¹⁴.




Figure 4: Variation in surgical treatment with age and route of presentation for breast cancers diagnosed in women in 2007

For symptomatic breast cancers, there was a significant relationship between surgical treatment and age at diagnosis; 90% of the women aged under 50 had surgical treatment recorded compared to only 74% of women aged 70-79 and 39% of patients aged 80 and

above. Significantly higher proportions of women with screen-detected breast cancer had surgery recorded in all age bands, and there was much less variation in surgical treatment in women under 80 years of age (Figure 4).

| Deprivation Quintile | Surgical Treatment | | | |
|----------------------|--------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | Total no. cases |
| Q1 Most deprived | 98% | 72% | 79% | 7344 |
| Q2 | 98% | 74% | 81% | 9201 |
| Q3 | 97% | 74% | 82% | 10582 |
| Q4 | 98% | 76% | 83% | 11385 |
| Q5 Least deprived | 98% | 78% | 85% | 11253 |
| All cases | 98% | 75% | 82% | 49765 |

Table 11: Variation in surgical treatment with deprivation quintile and route of presentation

For symptomatic breast cancers, there was a significant relationship between surgical treatment and deprivation quintile, with women in the most deprived group being less likely to have surgery (72% compared to 78% in women in the least

deprived group) (Table 11). This variation with deprivation was not apparent in women with screen-detected breast cancer, and significantly higher proportions of women with screen-detected breast cancer had surgery in all deprivation quintiles.

Surgical Treatment Outcomes

Breast Conserving Surgery and Mastectomy

In the UK as a whole, 57% of women with surgically treated breast cancer had breast conserving surgery and the remaining (43%) had a mastectomy. The mastectomy rate was significantly lower for screen-detected breast cancers (27% compared to 53% for symptomatic cancers), and a higher proportion of screen-detected breast cancers had breast conserving surgery in every age band (Figure 5). This difference is likely to be mainly due to the smaller size of the screen-detected cancers.

The proportions of women treated with breast conserving surgery were always significantly higher for screen-detected breast cancers in all deprivation quintiles (Table 12). For symptomatic breast cancers there is a statistically significant relationship between the proportion of cancers treated with breast conserving surgery and deprivation; the proportion of breast cancers treated with breast conserving surgery increased from 44% in the most deprived quintile to 50% in the least deprived quintile. This pattern is similar to that reported in other series such as Raine et al¹⁴ who analysed breast cancers diagnosed in women aged 50 years and above in 1999-2006. However, the more recent data in the present study




Figure 5: Variation in breast conserving surgery with age and presentation route for breast cancers diagnosed in women in 2007

show a narrowing of the variation with deprivation (1999/2006: 54% versus 64%; 2007: 44% versus 50%) suggesting that this inequity may be decreasing. The relationship between surgical procedure and deprivation is not apparent for screen-detected breast cancers.

| Deprivation Quintile | Breast Conserving Surgery | | | Total no. cases |
|----------------------|---------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 72% | 44% | 54% | 5797 |
| Q2 | 73% | 46% | 55% | 7419 |
| Q3 | 74% | 47% | 57% | 8635 |
| Q4 | 73% | 49% | 58% | 9478 |
| Q5 Least deprived | 74% | 50% | 59% | 9612 |
| All cases | 73% | 47% | 57% | 40941 |

Table 12: Variation in breast conserving surgery with deprivation quintile and route of presentation

Surgical Treatment Outcomes

The overall mastectomy rate for women with invasive breast cancer was significantly higher than that for women with non/micro-invasive breast cancer (44% compared to 38%). For symptomatic cases, the mastectomy rate for non/micro-invasive breast cancers was slightly but not significantly lower (51% compared to 53% for invasive breast cancers), while for screen-detected cases, the mastectomy rate for non/micro-invasive breast cancers was significantly higher (30% compared to 26% for invasive breast cancers). The latter may reflect the higher rate of immediate reconstruction for non/micro-invasive breast cancers which form a much higher proportion of screen-detected cases (20% compared to 8% of

symptomatic breast cancers). Overall, 10% of all mastectomies undertaken on female breast cancer patients in 2007 were for non-invasive disease. The pattern of care for non/micro-invasive breast cancers diagnosed through the NHS Breast Screening Programme highlights the difficulties in determining the optimal management for this disease, which can result in patients with a non-life threatening condition having more radical surgery than those with invasive breast cancer. Due to this uncertainty, the NICE Guideline *Early and locally advanced breast cancer: Diagnosis and treatment*¹⁵ recommends that all patients with screen-detected non-invasive breast cancer should be entered into the Sloane Project¹⁶.

Mastectomy: Breast Conserving Surgery Ratio

The national reference figure for the MX:BCS ratio has previously been reported as 1:1¹⁷, which assumes that the proportions of mastectomies (MX) and breast conserving surgery (BCS) operations are the same. In the UK as a whole for all breast cancers diagnosed in women, the MX:BCS ratio was 1:1.3. This is indicative of a higher level of BCS than anticipated by the national reference figure of 1:1. The MX:BCS ratio for all breast

cancers varied from 1:1.1 in the North West, Northern & Yorkshire and Trent regions to 1:1.7 in the Oxford region. The MX:BCS ratio was lower (i.e. the proportion of cases treated with BCS was higher) for screen-detected breast cancers than for symptomatic breast cancers (1:2.7 compared to 1:0.9), and for non/micro-invasive breast cancers than for invasive breast cancers (1:1.6 compared to 1:1.3).

| Age Band (years) | MX:BCS ratio | | |
|------------------|-----------------|-------------|---------|
| | Screen-detected | Symptomatic | Overall |
| <40 | - | 1:0.8 | 1:0.8 |
| 40-49 | 1:2.3 | 1:1.0 | 1:1.0 |
| 50-59 | 1:2.6 | 1:1.1 | 1:1.8 |
| 60-69 | 1:2.9 | 1:1.0 | 1:1.8 |
| 70-79 | 1:2.5 | 1:0.7 | 1:1.0 |
| 80+ | 1:1.5 | 1:0.6 | 1:0.7 |
| All ages | 1:2.7 | 1:0.9 | 1:1.3 |

| Age Band (years) | MX:BCS ratio | | |
|------------------|-----------------|-------------|---------|
| | Screen-detected | Symptomatic | Overall |
| <40 | - | 1:0.8 | 1:0.8 |
| 40-49 | 1:2.3 | 1:1.0 | 1:1.0 |
| 50-59 | 1:2.6 | 1:1.1 | 1:1.8 |
| 60-69 | 1:2.9 | 1:1.0 | 1:1.8 |
| 70-79 | 1:2.5 | 1:0.7 | 1:1.0 |
| 80+ | 1:1.5 | 1:0.6 | 1:0.7 |
| All ages | 1:2.7 | 1:0.9 | 1:1.3 |

Tables 13a and 13b: Variation in the MX:BCS ratio with a) age and route of presentation and b) deprivation quintile and route of presentation

Surgical Treatment Outcomes

MX:BCS ratios were also influenced by age and deprivation status. Because of the very high proportion of breast conserving surgery in women with screen-detected breast cancer, the overall MX:BCS ratio was lowest (1:1.8) in women aged 50-69 (Table 13a). For symptomatic breast cancers, the MX:BCS ratio was considerably higher in the youngest and oldest age bands (1:0.8 in women aged less than 40, 1:0.7 in women aged 70-79 and 1:0.6 in women aged 80 and over), indicating a relatively lower proportion of breast conserving surgery in these age bands. These age-related differences were less apparent in women with screen-detected breast cancers.

The overall MX:BCS ratio decreased with decreasing deprivation (from 1:1.2 in the most deprived quintile to 1:1.5 in the least deprived quintile), indicating a

higher proportion of breast conserving surgery in women in the least deprived quintile (Table 13b). When the deprivation data were spilt by route of presentation, a similar pattern was evident for symptomatic but not screen-detected breast cancers, and in every deprivation quintile the MX:BCS ratio was lower (i.e. the proportion of breast conserving surgery was higher) for screen-detected breast cancers.

The MX:BCS ratio increased from 1:5.1 and 1:2.6 for NPI Excellent and Good Prognostic Group breast cancers to 1:1.4 for Moderate Group 1 cancers, 1:0.9 for Moderate Group 2 cancers and 1:0.4 for Poor Prognosis Group cancers. These data are consistent with the proportion of mastectomy operations increasing with increasing invasive tumour size in the poorer prognostic groups.

Immediate Reconstruction

The time period covered in this report precedes the publication in 2009 of the NICE Guideline *Early and locally advanced breast cancer: Diagnosis and treatment*¹⁵ which states that immediate breast reconstruction should be discussed with all patients who are considering having a mastectomy, and offered as an option except where significant co-morbidity or (the need for) adjuvant therapy may preclude this choice.

In England, Wales and Scotland in 2007, 17,218 women with breast cancer were treated with mastectomy and

11% of these had an immediate reconstruction recorded. 10% of invasive breast cancers and 27% of non/micro-invasive breast cancers had immediate reconstruction recorded. Screen-detected cancers were more likely to have immediate reconstruction (16% compared to 10% of symptomatic cancers). This may be due, in part, to the higher proportion of screen-detected breast cancers that are non/micro-invasive (20% compared to 8% for symptomatic cases), and to the increased need for adjuvant therapy for symptomatic invasive breast cancers.

Surgical Treatment Outcomes

| Age Band (years) | Immediate Reconstruction | | | Total no. cases |
|------------------|--------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 26% | 26% | 1063 |
| 40-49 | 22% | 20% | 20% | 3102 |
| 50-59 | 24% | 12% | 17% | 3885 |
| 60-69 | 12% | 5% | 8% | 4216 |
| 70-79 | 4% | 1% | 1% | 3293 |
| 80+ | 0% | 0% | 0% | 1654 |
| All ages | 16% | 10% | 11% | 17213 |

| Deprivation Quintile | Immediate Reconstruction | | | Total no. cases |
|----------------------|--------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 12% | 10% | 10% | 2631 |
| Q2 | 13% | 10% | 10% | 3225 |
| Q3 | 16% | 10% | 11% | 3631 |
| Q4 | 19% | 10% | 12% | 3860 |
| Q5 Least deprived | 17% | 11% | 13% | 3834 |
| All cases | 16% | 10% | 11% | 17181 |

Tables 14a and 14b: Variation with a) age and route of presentation and b) deprivation quintile and route of presentation in the proportions of immediate reconstruction for breast cancers treated with mastectomy

Immediate reconstruction rates were also influenced by age and deprivation status (Tables 14a and 14b). Older women were less likely to have immediate reconstruction; no women aged 80 or over and only 1% of women aged 70 to 79 had immediate reconstruction, compared to 26% of women aged under 40. In women aged between 50 and 79 years, those with screen-detected breast cancer were significantly more likely to have an immediate reconstruction.

Among the women with screen-detected breast cancer treated with mastectomy, immediate reconstruction was significantly more likely to be undertaken in those from a less deprived background (12% of women in the most deprived quintile (Q1) compared with 19% and 17% in the least deprived quintiles (Q4 and Q5)). This relationship was not apparent for symptomatic breast cancers, but in all but the most deprived quintile (Q1), women with symptomatic breast cancer were significantly less likely to undergo immediate reconstruction than those with screen-detected breast cancer.

Figure 6 shows that 16% of the invasive breast cancers in the Excellent Prognostic Group (EPG) had an immediate reconstruction, compared to only 6% of the cancers in the Poor Prognosis Group (PPG). This reflects the increased frequency of the complicating factors associated with adjuvant therapy^{18,19} in women with breast cancers in the worse prognostic groups.




Figure 6: Immediate reconstruction after mastectomy for invasive breast cancers

Surgical Treatment Outcomes

Sentinel Lymph Node Biopsy (SLNB)

In England and Wales, 32% of women with surgically treated breast cancer had a SLNB recorded. This varied from 23% in the Trent region to 54% in the Oxford region. Of the cancers known to have received a SLNB, a significantly higher proportion of invasive breast cancers received a SLNB (35% compared to 13% of non/micro-invasive cancers). The use of SLNB has increased over time; in 2006 only 27% of invasive breast cancers had this procedure⁵.

The use of SLNB was higher for women with screen-detected breast cancer than for symptomatic women

(38% compared to 29%). More recent data from the NHSBSP demonstrate that 58% of screening patients had a SLNB in 2008/09. Over 80% of UK breast surgeons have now been trained in SLNB²⁰ so it is anticipated that the increased use of this technique, particularly in the treatment of early breast cancer, will continue. Women with invasive breast cancers in the Excellent Prognostic Group (EPG) were significantly more likely to undergo a SLNB; 50% of EPG cancers had a SLNB compared to 20% of cancers in the Poor Prognostic Group. This may in part explain the higher use of SLNB for screen-detected breast cancers.

| Age Band (years) | Sentinel lymph node biopsy | | | Total no. cases |
|------------------|----------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| <40 | - | 28% | 28% | 1760 |
| 40-49 | 34% | 31% | 31% | 5719 |
| 50-59 | 38% | 31% | 35% | 9709 |
| 60-69 | 38% | 30% | 35% | 10974 |
| 70-79 | 37% | 27% | 29% | 5852 |
| 80+ | 29% | 20% | 20% | 2535 |
| All ages | 38% | 29% | 32% | 36549 |

| Deprivation Quintile | Sentinel lymph node biopsy | | | Total no. cases |
|----------------------|----------------------------|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | |
| Q1 Most deprived | 37% | 26% | 30% | 5107 |
| Q2 | 36% | 26% | 30% | 6567 |
| Q3 | 37% | 28% | 31% | 7681 |
| Q4 | 37% | 29% | 32% | 8490 |
| Q5 Least deprived | 40% | 33% | 36% | 8627 |
| All cases | 38% | 29% | 32% | 36472 |

Tables 15a and 15b: Variation with a) age and route of presentation and b) deprivation quintile and route of presentation in the proportions of breast cancers having a sentinel lymph node biopsy

Older women were significantly less likely to have a SLNB; 20% of women aged at least 80 years had the procedure compared with 32% of all patients (Table 15a). Women in the least deprived quintile (Q5) were

significantly more likely to have a SLNB (36% compared to 30% in Q1 overall), regardless of whether their cancers were detected symptomatically or through screening (Table 15b).

Surgical Treatment Outcomes

Length of Hospital Stay

Length of hospital stay is recorded as the number of days from the date of admission to the date of discharge for the most radical surgical treatment. Data were available for England and Scotland only.

Of the 38,032 surgically treated female breast cancer cases diagnosed in England and Scotland, 21% had no length of stay recorded. This varied from 9.7% in the North West region to 30.6% in the Thames region. For the remaining 30,156 cases, 57% had a length of stay

of between 2 to 5 days. Women who had a mastectomy (M_x) stayed in hospital longer than women who had breast conserving surgery (BCS) (Figure 7). 70% of women treated with breast conserving surgery were discharged from hospital within 2 days compared with 21% of those treated with a mastectomy. The median number of days a woman treated with breast conserving surgery stayed in hospital was 2 days, compared to 4 days for mastectomy cases.




Figure 7: Variation in length of stay with surgical procedure for female breast cancer patients treated with breast conserving surgery (BCS) or mastectomy (M_x)

These overall differences were not affected by presentation route but, for women with symptomatic breast cancer, the median length of stay varied from 2 days in the West Midlands to 4 days in Scotland and the North West region. The median length of stay for patients with screen-detected breast cancer was 2 days and did not vary between regions. The broader spread of length

of stay observed for mastectomy patients is consistent with the increased risk of developing complications after mastectomy and reconstructive surgery. The shorter overall length of stay for patients with screen-detected cancer reflects the higher breast conserving surgery rate for patients with these better prognosis cancers (73% compared to 47% for symptomatic cancers).




Figure 8: Variation in length of stay with age band for female breast cancer patients

Surgical Treatment Outcomes

Length of stay varied significantly with age at diagnosis (Figure 8). Within the 50-69 year age band, 28% of women remained in hospital for 1 day or less, with a further 24% staying in hospital for 2 days. In contrast, 27% of women aged 80 or over spent between 6 and 10 days in hospital following their surgery compared to only 15% of women in the 50-69 year age band.

Women aged 70 years and over formed 63% of the 72 patients who remained in hospital for 20 days or longer. Only 42% of women aged less than 50 stayed in hospital for 2 days or less compared with 52% of women aged 50-69 years; this may reflect the higher mastectomy with immediate reconstruction rates in the younger women.




Figure 9: Variation in length of stay with deprivation quintile for female breast cancer patients

Length of stay also varied significantly with deprivation quintile (Figure 9). When compared to women in the least deprived quintile, women in the most deprived quintile were less likely to stay in hospital for 2 days or less (42% compared with 48%). They were also more likely to stay in

hospital for more than 5 days (23% compared to 17% for women in the least deprived quintile). As breast units implement 23 hour mastectomies, it will be interesting to monitor these data to ascertain which patients are able to access this emerging model of care.

Repeat Operation Rates

In England, Wales and Scotland significantly more women with non/micro-invasive breast cancers were known to have had a repeat operation (25% compared to 14% for women with invasive breast cancer). The overall repeat operation rate for women presenting via screening was higher than in those presenting symptomatically (18% versus 15%). This reflects the higher proportion of non/micro-invasive cancers and the greater use of breast conserving surgery in women with screen-detected breast cancer.

For both screen-detected and symptomatic cases, repeat operation rates varied with the age of the woman; older women were less likely to have repeat operations (Table 16). The trend was more obvious for symptomatic women; only 15% of the surgically treated cancers in women aged 70 or over had a repeat operation, compared to 20% of those aged less than

40. There was no relationship between repeat operation rates and deprivation for women with screen-detected or symptomatic breast cancers.

| Age Band (years) | Screen-Detected | | Symptomatic | |
|------------------|------------------|-----------------|-------------|-----------------|
| | Repeat operation | Total no. cases | Repeat | Total no. cases |
| <40 | - | - | 20% | 1696 |
| 40-49 | 19% | 155 | 19% | 5358 |
| 50-59 | 19% | 5235 | 17% | 4125 |
| 60-69 | 17% | 6714 | 15% | 4042 |
| 70-79 | 17% | 1315 | 10% | 4605 |
| 80+ | 13% | 79 | 5% | 2399 |
| All ages | 18% | 13503 | 15% | 22225 |

Table 16: Variation in repeat operation rate with age for women with screen-detected and symptomatic breast cancers

Surgical Treatment Outcomes

30 Day Mortality

In 2007, 860 women (2%) with breast cancer died within 30 days of their diagnosis. 51% of these were death certificate only (DCO) registrations where the date of diagnosis was the date of death. 87% of the DCO registrations were aged 70 years or more; 58 were below 70 years of age. 78% of the 426 non-DCO cases who died within 30 days of diagnosis were aged 70 years or more; 94 were below 70 years of age. Only 7 of the non-DCO cases who died within 30 days had undergone surgical treatment for their breast cancer.

Of the DCO cases, 66% died from breast cancer, 4% from other cancers and 30% from other non-cancer or unknown causes. Of the non-DCO cases who died within 30 days, 64% died of breast cancer. The latter varied with age at diagnosis; 72% of those aged less than 70 died of breast cancer compared with 62% of

those aged 70 years and over. Overall, 9% of the non-DCO cases died from another cancer and 27% died from other causes. The proportion dying of another cancer was higher in women aged less than 70 years (12% compared to 8% in women aged 70 and over) and the proportion dying from other causes was lower (12% compared to 27% in women aged 70 and over).

Deprivation status was not known for 7 women who died within 30 days of their diagnosis of breast cancer. There was a significant relationship between 30 day mortality and deprivation; 2.8% of all cases and 1.4% of non-DCO cases dying within 30 days were women in the most deprived quintile compared with only 1.4% and 0.7% respectively of women in the least deprived quintile (Table 17).

| Deprivation quintile | All cases | Total no. cases | Non-DCO cases | Total no. cases |
|----------------------|-----------|-----------------|---------------|-----------------|
| Q1 Most deprived | 2.8% | 165 | 1.4% | 84 |
| Q2 | 2.4% | 180 | 1.2% | 88 |
| Q3 | 2.2% | 192 | 1.1% | 93 |
| Q4 | 1.9% | 185 | 1.0% | 95 |
| Q5 Least deprived | 1.4% | 131 | 0.7% | 62 |
| All cases | 2.1% | 853 | 1.0% | 422 |

Table 17: Variation with deprivation quintile in the proportion of all breast cancer patients and DCO patients who died within 30 days of their diagnosis

Surgical Treatment Outcomes

Surgical Caseload

In England, Wales and Northern Ireland in 2007, 524 surgeons were recorded as having treated a total of 36,695 women with breast cancer. Their caseload ranged from 1 to 258 cases, with a median caseload of 69. 385 (73%) of the surgeons treated at least 30 breast

cancer cases, and therefore met or exceeded the recommended caseload. Figure 10 shows the mastectomy rate for the 524 surgeons. 27 surgeons have a significantly high mastectomy rate and 44 surgeons have a significantly low mastectomy rate.




Figure 10: Variation in mastectomy rates between surgeons

Key Findings

- Survival analyses were performed on 39,879 invasive and micro-invasive breast cancers diagnosed in women in the UK between 1 April 2002 and 31 March 2003, and 44,069 invasive and micro-invasive breast cancers diagnosed between 1 January 2007 and 31 December 2007.
- 1-year relative survival was significantly higher for women diagnosed with breast cancer in 2007 (96% compared to 94% in 2002/03).
- 1-year and 5-year relative survival rates were significantly higher for women with screen-detected breast cancer (100% compared to 93%-94% at 1 year and 97% compared to 77% at 5 years).
- For women with symptomatic breast cancer, 5-year relative survival decreased with age (from 86% in women aged 40-49 years to only 62% in women aged 80 years and above).
- For women with screen-detected breast cancer, 1-year relative survival was approximately 100% for all age bands, indicating that these women were no more likely to die than those in the population as a whole.
- 1-year and 5-year relative survival rates for women with symptomatic breast cancer were significantly lower than those for women with screen-detected breast cancer in all deprivation quintiles.
- For women with symptomatic breast cancer, there were marked decreases in 1-year survival with deprivation (from 90%-92% for women in the most deprived quintile to 96%-97% for women in the least deprived quintile). These differences were not apparent for women with screen-detected breast cancer.
- For women with screen-detected breast cancer, 5-year relative survival increased from 94% in the most deprived quintile to 99% in the least deprived quintile. There was a much more marked difference in 5-year relative survival between the most and least deprived quintiles for women with symptomatic breast cancer; 5-year relative survival being 68% in the most deprived quintile and 83% in the least deprived quintile.

Relative Survival

Survival analyses were performed on 39,879 invasive and micro-invasive breast cancers diagnosed in women in the UK between 1 April 2002 and 31 March 2003, and 44,069 invasive and micro-invasive breast cancers diagnosed between 1 January 2007 and 31 December 2007. Relative survival can be interpreted as the ratio

between the survival in the patient group examined and that in the general population. If the relative survival rate is 100%, the survival of the patient group is the same as that of the general population. The proportion of patients dying of breast cancer was not determined.

Mortality

The proportions of women in both cohorts who were alive on 31 December 2008 are shown in Table 18. Cases with unknown screening status have been excluded from this table. In the 2002/03 cohort, 70% of women were recorded as being alive on 31 December 2008, and in the 2007 cohort, 84% were

alive on 31 December 2008. These percentages give no indication of the reason for the deaths. Notably, in both cohorts, women with symptomatic breast cancer were more likely to have died than women in the same age band with screen-detected breast cancer.

| Age Band (years) | 2002/03 Cohort Alive on 31 December 2008 | | | | 2007 Cohort Alive on 31 December 2008 | | | |
|------------------|---|-------------|-----------|-----------------|--|-------------|-----------|-----------------|
| | Screen-detected | Symptomatic | All cases | Total no. cases | Screen-detected | Symptomatic | All cases | Total no. cases |
| <40 | - | 78% | 78% | 2305 | - | 91% | 91% | 2007 |
| 40-49 | 93% | 83% | 83% | 6090 | 96% | 93% | 93% | 6491 |
| 50-59 | 93% | 79% | 86% | 12069 | 98% | 88% | 93% | 9997 |
| 60-69 | 91% | 69% | 79% | 9493 | 97% | 84% | 91% | 11446 |
| 70-79 | 86% | 55% | 57% | 7765 | 94% | 77% | 79% | 7804 |
| 80+ | 83% | 26% | 26% | 6436 | 88% | 53% | 54% | 6762 |
| All ages | 92% | 63% | 70% | 44158 | 97% | 79% | 84% | 44507 |

Table 18: Variation with age and route of presentation in the proportion of women diagnosed with breast cancers in 2002/03 and 2007 known to be alive on 31 December 2008

1-Year and 5-Year Relative Survival

5-year relative survival for all women diagnosed with invasive or micro-invasive breast cancer in 2002/03 was 81% (Table 19). 5-year relative survival for women with screen-detected breast cancer was significantly higher than that for women with symptomatic breast cancer (97% compared to 77%). 1-year relative survival was significantly higher for

women diagnosed with breast cancer in 2007 (96% compared to 94% in 2002/03). For women with screen-detected breast cancer 1-year relative survival was 100% in both cohorts. 1-year relative survival for women with symptomatic breast cancer was significantly lower; 93% in the 2002/03 cohort and 94% in the 2007 cohort.

| Presentation route | 2002/03 Cohort | | 2007 Cohort |
|--------------------|------------------------------------|------------------------------------|------------------------------------|
| | 1-year relative survival (LCI-UCI) | 5-year relative survival (LCI-UCI) | 1-year relative survival (LCI-UCI) |
| Screen-detected | 100.0% (99.8%-100.2%) | 97.1%(96.5%-97.6%) | 100.2% (100.1%-100.4%) |
| Symptomatic | 92.8% (92.4%-93.1%) | 76.8% (76.2%-77.4%) | 94.0% (93.6%-94.3%) |
| All cases | 94.3% (94.1%-94.6%) | 81.3% (80.8%-81.8%) | 95.8% (95.6%-96.0%) |

Table 19: 1-year and 5-year relative survival rates with presentation route for women diagnosed with invasive/micro-invasive breast cancer in 2002/03 and 2007 (LCI = 95% lower confidence interval; UCI = 95% upper confidence interval)

Survival

Age

Figure 11 shows how 5-year relative survival varied with age at diagnosis for women diagnosed with invasive or micro-invasive breast cancer in 2002/03. For women with screen-detected breast cancer, there was little variation in 5-year relative survival with age. At 104%, 5-year relative survival in women aged 70-79, was higher than in the general population, but this difference was not statistically significant. 5-year relative survival was significantly lower in every age band for women with symptomatic breast cancer. For women aged 40 and above with symptomatic cancer, there was a marked decrease in 5-year relative survival with age; 5-year relative survival decreased from 86% in women aged 40-49 years to only 62% in women aged 80 years and above. 5-year relative survival for women aged less than 40 years was lower than that for women aged 40-49 and 50-59 years.

1-year relative survival was similar in both cohorts of women with screen-detected breast cancer, and did not vary significantly with age band in either cohort (Table 20). 1-year relative survival was approximately 100% all age bands, indicating that women with screen-detected breast cancer were no more likely to die than those in the population as a whole. With the exception of women in the 40-49 year age band in 2007, 1-year relative survival for women with




Figure 11: 5-year relative survival with age and presentation route for women diagnosed with invasive/micro-invasive breast cancer in 2002/03

symptomatic breast cancer was significantly lower than that for women with screen-detected breast cancer. For women with symptomatic breast cancer, there were marked decreases in 1-year survival with age (from around 98% in women aged 40-49 years to 86% in women aged 80 years and above). 1-year relative survival rates were generally slightly higher in the 2007 cohort, and these differences were statistically significant in women aged 60 years and over.

| Age Band (years) | 1-year relative survival Screen-detected (LCI-UCI) | | 1-year relative survival Symptomatic (LCI-UCI) | |
|------------------|--|------------------------|--|---------------------|
| | 2002/03 Cohort | 2007 Cohort | 2002/03 Cohort | 2007 Cohort |
| <40 | - | - | 96.9% (96.1%-97.6%) | 97.9% (97.2%-98.5%) |
| 40-49 | 100.3% (100.3%-100.3%) | 99.4% (94.2%-100.1%) | 98.1% (97.7%-98.5%) | 98.1% (97.7%-98.4%) |
| 50-59 | 99.9% (99.6%-100.0%) | 100.1% (99.9%-100.2%) | 96.8% (96.3%-97.3%) | 97.0% (96.5%-97.5%) |
| 60-69 | 100.1% (99.7%-100.4%) | 100.3% (100.0%-100.4%) | 93.6% (92.9%-94.3%) | 94.8% (94.1%-95.5%) |
| 70-79 | 101.0% (99.5%-101.8%) | 100.9% (100.0%-101.4%) | 90.5% (89.7%-91.3%) | 92.6% (91.8%-93.3%) |
| 80+ | 103.4% (85.7%-106.2%) | 102.2% (93.5%-105.6%) | 83.0% (81.7%-84.2%) | 86.1% (85.0%-87.2%) |

Table 20: 1-year relative survival rates with age and presentation route for women diagnosed with invasive/micro-invasive breast cancer in 2002/03 and 2007 (LCI = 95% lower confidence interval; UCI = 95% upper confidence interval)

The relatively low survival rate for older patients is partly because of the high proportion of cases without surgery in this group. Figure 12 shows that the relative survival rate for patients without surgery is significantly lower than for those with surgery.

Deprivation

For women with screen-detected breast cancer, 5-year relative survival increased by 5% points from 94% in the most deprived quintile (Q1) to 99% in the least deprived quintile (Q5) (Figure 13), but this increase was not statistically significant. 5-year relative survival was significantly lower in every deprivation quintile for women with symptomatic breast cancer, and there was a much more marked and statistically significant 15% points difference between the 5-year relative survival of women in the most deprived quintile (Q1, 68%) and in the least deprived quintile (Q5, 83%).

1-year relative survival was similar for women diagnosed with screen-detected breast cancer in 2002/03 and 2007, and did not vary significantly with deprivation quintile in either cohort (Table 21). 1-year relative survival was approximately 100% in all deprivation quintiles, indicating that women with screen-detected breast cancer were no more likely to die than those in the population as a whole. 1-year relative survival for women with symptomatic breast cancer was significantly lower than that for women with screen-detected breast cancer in all deprivation quintiles in both cohorts. For women with symptomatic breast cancer, there were marked variations in 1-year survival with deprivation (from 90%-92% for women in the most deprived quintile (Q1) to 96%-97% for women in the least deprived quintile (Q5)). 1-year relative survival was generally slightly higher in the 2007 cohort, but the differences were not statistically significant.




Figure 12: Variation in 1-year relative survival rate with age for symptomatic invasive and micro-invasive breast cancers diagnosed in 2007




Figure 13: Variation in 5-year relative survival with deprivation quintile and presentation rate for women diagnosed with invasive/micro-invasive breast cancer in 2002/03

| Deprivation Quintile | 1-year relative survival Screen-detected (LCI-UCI) | | 1-year relative survival Symptomatic (LCI-UCI) | |
|----------------------|---|---------------------------|---|------------------------|
| | 2002/03 Cohort | 2007 Cohort | 2002/03 Cohort | 2007 Cohort |
| Q1 Most deprived | 99.4% (98.6%-99.9%) | 99.8% (99.2%-100.1%) | 89.6% (88.6%-90.5%) | 91.2% (90.3%-92.0%) |
| Q2 | 99.6% (99.0%-100.0%) | 100.0% (99.5%-100.3%) | 91.1% (90.2%-91.9%) | 92.8% (92.0%-93.5%) |
| Q3 | 100.0% (99.5%-100.3%) | 100.0% (99.6%-100.3%) | 92.9% (92.1%-93.6%) | 93.7% (92.9%-94.3%) |
| Q4 | 100.5% (100.1%-100.7%) | 100.6% (100.3%-100.8%) | 93.9% (93.2%-94.6%) | 94.8% (94.1%-95.4%) |
| Q5 Least deprived | 100.4% (99.9%-100.6%) | 100.6% (100.2%-100.8%) | 95.6% (95.0%-96.2%) | 96.5% (95.9%-97.1%) |

Table 21: 1-year relative survival rates with deprivation quintile and presentation route for women diagnosed with invasive/micro-invasive breast cancer in 2002/03 and 2007
(LCI = 95% lower confidence interval; UCI = 95% upper confidence interval)

References

1. Department of Health (2010). Equity and excellence: Liberating the NHS. July 2010
2. Department of Health (2010). Liberating the NHS: Transparency in Outcomes – a framework for the NHS, July 2010
3. National Institute for Health and Clinical Excellence (2011). Quality Standards for Breast Cancer <http://www.nice.org.uk/guidance/qualitystandards/indevelopment/breastcancer.jsp>
4. National Cancer Action Team (2010). National Cancer Peer Review Programme Evidence Guide for: Breast MDT, version 2. February 2010
5. NHS Cancer Screening Programme and NCIN (2009). All Breast Cancer Report, October 2009
6. NHS Connecting for Health. Data Standards: Cancer Registration Data Set. <http://www.isb.nhs.uk/documents/isb-0056/dscn-09-2005> June 2005
7. NHS Connecting for Health. Hospital Episodes Statistics www.connectingforhealth.nhs.uk/systemsandservices/sus/delivery/hes. Date accessed 17.03.11
8. Communities and Local Government (2008). The English Indices of Deprivation 2007, Communities and Local Government: London. March 2008
9. Chiu, L.F. (2003). Inequalities of Access to Cancer Screening: A Literature Review. Cancer Screening Series No 1 December 2003
10. Galea, M.H., Blamey, R.W., Elston, C.E., Ellis, I.O. (1992). The Nottingham Prognostic Index in primary breast cancer. *Breast Cancer Research Treatment*, 1992, 22: 207–219.
11. Slamon, D.J., Clark, G.M., Wong, S.G., Levin, W.J., Ullrich, A., McGuire, W.L, (1987). Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science* 1987;**235**(4785):177-82.
12. Vogel, U. (2010). Confirmation of a low HER2 positivity rate of breast carcinomas - limitations of immunohistochemistry and in situ hybridization. *Diagn Pathol*. 2010 Jul 29;5:50.
13. NHSBSP (2010). An audit of screen-detected breast cancers for the year of screening April 2008 to March 2009. NHS Breast Screening Programme, May 2010
14. Raine, R., Wong, W., Scholes, S., Ashton, C., Obichere, A., Ambler, G. (2010). Social variations in access to hospital care for patients with colorectal, breast, and lung cancer between 1999 and 2006: retrospective analysis of hospital episode statistics. *BMJ*. 2010 Jan 14;340.
15. National Institute for Health and Clinical Excellence. (2009) Breast cancer (early and locally advanced): diagnosis and treatment. Clinical Guideline 80; February 2009
16. The Sloane Project www.sloaneproject.org.uk
17. Department of Health (2009). Cancer Commissioning Guidance, amended. December 2009
18. Vandeweyer, E., Deraemaeker, R., Nogaret, J.M., Hertens, D. (2003) Immediate Breast Reconstruction with Implants and Adjuvant Chemotherapy: A Good Option? *Acta chir belg*, 2003, 103, 98-101
19. National Mastectomy and Breast Reconstruction Audit (2010) A national audit of provision and outcomes of mastectomy and breast reconstruction surgery for women in England. Third Annual Report. June 2010
20. Breast Educational and Training portfolio RCS England NEW START SLNB training programme; <http://www.baso.org/content/AbS-RCS.asp> Date accessed 17.03.11

Appendix 1 - Participating Surgeons

| | | | |
|----------------------|-----------------------|--------------------|--------------------|
| Mr TI Abdullah | Miss J Bonnema | Mr MA Chaudary | Miss SE Downey |
| Mr R Achuthan | Mr R Bonomi | Mr IA Cheema | Mr EH Drabble |
| Mr JK Adjogatse | Mr R Bourne | Mr KL Cheung | Prof PJ Drew |
| Miss AMG Aertssen | Mr HA Bradpiece | Mr CI Chianakwalam | Miss PV Dudani |
| Mr AC Agombar | Mr PA Braithwaite | Mr KH Chin | Mr TJ Duffy |
| Mr A Agrawal | Miss MD Bramley | Mr AK Chouhan | Miss EJ Duggan |
| Mr IA Ahmed | Miss RM Bright-Thomas | Mr KR Clark | Miss JM Dunn |
| Mr MSM Al-Dubaisi | Mr JB Bristol | Miss PJ Clarke | Miss P Durning |
| Mr MAA Al-Gailani | Mr PT Brookes | Mr D Clarke | Mr GW Dyke |
| Mr D Ali | Mr R Brookstein | Mr DJ Clarke | Mr S Dzumhur |
| Mr SM Allan | Mr DA Browell | Mr PR Clothier | Mr SR Ebbs |
| Mr WH Allum | Mr IM Brown | Mr RA Cochrane | Mr S Ellenbogen |
| Miss A Anand | Mr JH Brown | Mr NJ Coombs | Mr DJ Ellis |
| Mr IMF Anwar | Mr R Brown | Mr MJ Cooper | Mr DW England |
| Mr TJ Archer | Mr RM Bryan | Mr GP Copeland | Prof OF Eremin |
| Mr PR Armitstead | Mr TE Bucknall | Mr AP Corder | Miss AA Evans |
| Mr AM Armstrong | Prof NJ Bundred | Miss CA Courtney | Mr NW Everson |
| Miss AC Athow | Mr GJ Byrne | Mr SP Courtney | Prof IS Fentiman |
| Mr RA Audisio | Mr MJ Callam | Mr E Coveney | Mr DJ Ferguson |
| Mr A Aukland | Mr JK Campbell | Mr PW Crane | Mr DRA Finch |
| Miss IAF Azmy | Mr PJ Cant | Mr DJ Crawford | Mr P Forouhi |
| Mr AD Baildam | Miss AR Carmichael | Mr RS Cummins | Mr KD Fortes-Mayer |
| Mr B Balasubramanian | Mr R Carpenter | Mr GH Cunnick | Miss CA Fowler |
| Mr D Banerjee | Mr M Carr | Mr RA Daoud | Mr JN Fox |
| Mr MM Barkeji | Mr NJ Carty | Mr TI Davidson | Miss A Francis |
| Mr LC Barr | Mr WG Case | Mr CJ Davies | Mr PB Frecker |
| Mr N Beechey-Newman | Mr SJ Cawthorn | Mr JW Dawson | Mr MH Galea |
| Miss AM Bello | Mr DR Chadwick | Mr MN Dehalvi | Mr NC Gallegos |
| Mr JR Benson | Miss LS Chagla | Mr AZ Demian | Mr A Gandhi |
| Mr DA Berstock | Mr HY Chan | Mr AJ Desai | Mr TCS Gate |
| Mr PK Bhaskar | Miss VPP Chandran | Mr E Dinakara Babu | Mr CA Gateley |
| Mr TS Bhatti | Mr S Chandrasekharan | Mr LA Donaldson | Miss JM Gattuso |
| Mr HM Bishop | Mr MV Chandrashekar | Miss MA Donlon | Mr R Gendy |
| Mr RD Bliss | Mr MJB Chare | Miss JS Donnelly | Mr PP George |
| Mr GP Boland | Mr GHGM Charfare | Mr PK Donnelly | Miss MW Ghilchik |

Appendix 1 - Participating Surgeons

| | | | |
|------------------------|------------------|--------------------|-----------------------|
| Mr S Ghosh | Mr PA Holland | Mr RJ Kennedy | Mr DM Matheson |
| Mr S Ghosh | Mr HW Holliday | Miss FS Kenny | Mr WA Maxwell |
| Mr JA Gill | Mr S Holt | Mr SN Khan | Mr FP McGinn |
| Mr DA Gilpin | Mr SDH Holt | Mr AR Khawaja | Mr AJ McLaren |
| Glenfield Hospital | Mr HMR Hoque | Mr NI Khonji | Miss PL McManus |
| Mr AB Gordon | Mr KM Horgan | Mr RM Kirby | Mr GAD McPherson |
| Mr PG Gough | Mr MMI Hussien | Mr SJ Kirk | Mr M Menon |
| Mr S Goyal | Mr IF Hutchinson | Mr P Kiruparan | Mr M Mirza |
| Mr MD Graham | Miss J Iddon | Mr MW Kissin | Mr VK Modgill |
| Miss AE Gray | Mr HT Ingle | Mr AM Klidjian | Mr AK Modi |
| Mr MHA Green | Mr ERI Inwang | Mr RA Knox | Prof K Mokbel |
| Mr MJ Greenall | Miss S Iqbal | Mr SR Kohlhardt | Mr IJ Monypenny |
| Mr CDM Griffith | Mr B Isgar | Mr JS Kokan | Miss CJD Mortimer |
| Mr AB Griffiths | Mr WWM Ismail | Mr KM Kolar | Mr AIH Mostafa |
| Mr NJ Griffiths | Mr LR Jackson | Mr TG Kovacs | Mr VP Murali Krishnan |
| Mr JW Groome | Mr V Jaffe | Mr IJ Laidlaw | Mr SM Musa |
| Mr E Gross | Mr MA Jahan | Mr MRJ Lansdown | Mr R Nangalia |
| Mr G Gui | Miss SC Jenkins | Miss SAM Laws | Mr A Nejim |
| Mr DJ Hadjiminias | Mr AJ Jewkes | Prof GT Layer | Mr MH Niayesh |
| Mr MT Hallissey | Mr JA Jibril | Mr MJR Lee | Mr S Nicholson |
| Mr HHA Hamed | Mr SARJB Jmor | Prof TWJ Lennard | Mr AJ Ogedegbe |
| Miss C Harding-MacKean | Mr JN Johnson | Mr RA Linforth | Mr AA Ojo |
| Mr JS Harisha | Mr OO Johnson | Mr JM Lund | Miss JL Ooi |
| Mr RNL Harland | Mr RC Johnson | Mr M Lwin | Mr MH Ornstein |
| Mr SA Harries | Mr AK Johri | Mr RD MacMillan | Mr JA Pain |
| Mr KP Harris | Miss C Jones | Miss FA MacNeill | Mr SJ Pain |
| Mr ME Harron | Miss LE Jones | Mr PR Maddox | Mr SJ Parker |
| Miss DA Hassanally | Miss SE Jones | Mr TK Mahapatra | Mr JR Parmar |
| Miss M Heitmann | Mr BG Jones | Mr JB Mancey-Jones | Mr R Parmeshwar |
| Mr C Hennessy | Mr LS Jones | Prof RE Mansel | Mr AR Patel |
| Mr MJ Higgs | Mr PA Jones | Mr JG Maroof | Mr AG Paterson |
| Mr CP Hinton | Prof AK Kakkar | Mr SK Marsh | Mr IS Paterson |
| Miss FJ Hoar | Mr B Kald | Mr SG Marshall | Mr ALG Peel |
| Miss RKF Hogben | Mr NHK Kazzazi | Mr L Martin | Mr G Peley |
| Mr C Holcombe | Mr CJ Kelley | Miss MP Matey | Mr JH Pereira |

Appendix 1 - Participating Surgeons

| | | | |
|--------------------------|---------------------|--------------------|--------------------|
| Mr D Perry | Miss C Roshan Lall | Miss SA Sloan | Mr R Vashisht |
| Miss LH Phipp | Mr ND Rothnie | Mr BM Smith | Miss R Vidya |
| Mr MR Pittam | Mr GT Royle | Mr SGT Smith | Mr RK Vijayakumar |
| Mr AK Poddar | Mr PC Runchman | Mr SJ Snooks | Mr R Vijh |
| Miss CJ Pogson | Miss JE Rusby | Mr RG Souter | Miss LS Vishwanath |
| Miss V Pope | Mr PC Rutter | Mr A Stacey-Clear | Miss AJ Waghorn |
| Mr MJ Pospiech | Miss ZA Sa'Ad | Miss MA Stebbing | Miss Y Wahedna |
| Mr R Prasad | Mr AK Sahu | Mr RD Stewart | Miss J Walls |
| Miss OD Predolac | Mr ZAS Saidan | Mr WJ Stirling | Mr RM Watkins |
| Mr CJ Pritchett | Mr JRC Sainsbury | Mr PS Stonelake | Mr RJ Watson |
| Mr AM Przyczyna | Mr AK Salih | Miss AT Stotter | Miss BE Weber |
| Mr JV Psaila | Mr AR Salman | Mr PN Strauss | Mr SM Whitehead |
| Mr NJ Purser | Mr AM Sammon | Mr AP Subash Kumar | Mr AJ Wilkinson |
| Prof AD Purushotham | Mr WA Samra | Mr RJ Sutton | Mr EV Williams |
| Mr G Querci della Rovere | Mr DGD Sandilands | Mr FM Swe | Mr MR Williams |
| Mr RM Rainsbury | Prof PD Sauven | Miss HM Sweetland | Mr NJ Williams |
| Mr NI Ramus | Mr SN Selvachandran | Miss AP Tansley | Mr RJL Williams |
| Mr AAH Rashed | Miss BN Shah | Miss N Tayal | Mr R Windle |
| Mr MD Rashed | Miss EF Shah | Miss AJ Taylor | Mr JHR Winstanley |
| Mr AH Rateme | Mr YM Sharaiha | Mr AR Taylor | Miss ZE Winters |
| Mr D Ravichandran | Mr AK Sharma | Mr JL Taylor | Mr MH Wise |
| Mr SSB Ravindra Bose | Mr NK Sharma | Mr KK Thakur | Mr GC Wishart |
| Mr O Ravisekar | Miss EJ Sharp | Mr SD Thomson | Mr JJ Wood |
| Mr SP Raymond | Miss KC Shenton | Miss AL Thorne | Miss L Wyld |
| Mr Z Rayter | Mr SG Shering | Mr S Thrush | Mr A Yelland |
| Miss EJ Redmond | Mr MS Shrotri | Mr JC Tresadern | Mr C Yiangou |
| Prof MWR Reed | Miss S Shrotria | Mr EPL Turton | Mr MMKI Youssef |
| Mr RA Reichert | Mr DM Sibbering | Mr HN Umeh | Mr C Zammit |
| Mr DA Rew | Mr SM Singh | Mr HC Umpleby | Miss K Zechmeister |
| Mr JV Roberts | Mr H Singhal | Mr T Usman | Surgeons in the |
| Miss NA Roche | Mr PS Sinha | Miss K Valassiadou | Scottish Cancer |
| Miss CR Rogers | Prof HD Sinnett | Mr D Valerio | Networks |
| Mr CA Rogers | Mr AI Skene | Mr S Varadarajan | |

Appendix 2 – Overall Data Quality and Completeness

Table A: Data items in each country included in the analyses

| Data item | England | Northern Ireland | Scotland | Wales |
|--|---------|------------------|----------|-------|
| Invasive size, invasive grade, nodal status and Nottingham Prognostic Index for invasive cancers | ✓ | ✓ | ✓ | ✓ |
| Vascular invasion | ✓ | | | ✓ |
| Oestrogen receptor status | ✓ | | ✓ | ✓ |
| Human epidermal growth factor receptor status | ✓ | | | ✓ |
| Surgery | ✓ | ✓ | ✓ | ✓ |
| Final therapeutic operation type | ✓ | ✓ | ✓ | ✓ |
| Sentinel lymph node biopsy | ✓ | | | ✓ |
| Immediate reconstruction | ✓ | | ✓ | ✓ |
| Length of stay | ✓ | | ✓ | |
| Number of operations | ✓ | | ✓ | ✓ |
| Surgical caseload | ✓ | ✓ | | ✓ |
| Survival | ✓ | ✓ | ✓ | ✓ |

Table B: Numbers and proportions of data items with known values (UK female and male)

| Data item | Number of cases | UK | England |
|--|-----------------|------|---------|
| UK cohort | 50,286 | - | - |
| Deprivation score | 50,088 | 100% | 100% |
| Ethnicity | 33,648 | 67% | 80% |
| Surgical treatment (BCS, MX and no surgery) | 43,573 | 87% | 86% |
| UK invasive cancers cohort | 44,782 | - | - |
| Surgical treatment (BCS, MX and no surgery) | 38,373 | 86% | 85% |
| UK invasive cancers cohort (BCS and MX only) | 36,182 | - | - |
| Invasive tumour size | 32,296 | 89% | - |
| Invasive tumour grade | 35,019 | 97% | - |
| Nodal status | 28,786 | 80% | - |
| NPI | 27,581 | 76% | - |
| Oestrogen receptor status | 21,553 | 60% | - |
| Progesterone receptor status | 14,582 | 40% | - |
| HER2 status | 15,874 | 44% | - |
| Sentinel lymph node biopsy | 16,809 | 46% | - |
| Immediate reconstruction | 7,168 | 20% | - |
| Length of stay | 27,159 | 75% | - |

Appendix 2 – Overall Data Quality and Completeness

Table C: Proportion of cases with unknown values

| Data Item | Screen-detected | Symptomatic |
|-----------------------------|-----------------|-------------|
| Age | 0% | 0% |
| Ethnicity | 30% | 34% |
| Deprivation | <1% | <1% |
| Surgical treatment | <1% | 19% |
| Invasive size | <1% | 16% |
| Invasive grade | <1% | 5% |
| Nodal status | 2% | 30% |
| Nottingham Prognostic Index | 3% | 34% |

Appendix 3 – Tumour Characteristics Data Completeness

Table D: Data completeness for invasive breast cancers diagnosed in females in 2007 (expressed as % of all invasive tumours)

| English region/ Celtic country | Number of invasive tumours | % with known invasive tumour size | % with known invasive tumour grade | % with known nodal status | % with known NPI | % with known vascular invasion status | % with known ER status | % with known PgR status | % with known HER2 status |
|-----------------------------------|----------------------------|-----------------------------------|------------------------------------|---------------------------|------------------|---------------------------------------|------------------------|-------------------------|--------------------------|
| Eastern | 4,194 | 90.0% | 90.7% | 78.6% | 76.6% | 16.2% | 46.0% | 28.5% | 39.3% |
| North West | 4,881 | 77.4% | 88.8% | 74.1% | 72.0% | 11.6% | 40.9% | 40.5% | 34.4% |
| Northern & Yorkshire | 4,824 | 88.1% | 92.7% | 53.6% | 53.1% | 27.3% | 60.7% | 53.1% | 48.1% |
| Oxford | 1,993 | 76.1% | 94.1% | 83.5% | 72.8% | 83.5% | 43.0% | 42.7% | 36.6% |
| South West | 5,949 | 78.5% | 88.6% | 74.1% | 72.0% | 72.0% | 51.9% | 38.3% | 38.0% |
| Thames | 7,521 | 70.8% | 86.7% | 50.6% | 45.7% | 8.5% | 31.8% | 29.7% | 25.9% |
| Trent | 3,501 | 41.0% | 87.4% | 40.6% | 39.7% | 11.4% | 43.9% | 11.5% | 38.8% |
| W Midlands | 4,225 | 84.5% | 92.5% | 80.6% | 79.1% | 79.9% | 96.1% | 86.8% | 87.0% |
| England | 37,088 | 76.4% | 89.7% | 65.3% | 62.5% | 34.8% | 50.7% | 40.9% | 42.1% |
| N Ireland | 1,132 | 84.5% | 85.2% | 84.3% | 75.6% | 0% | 0% | 0% | 0% |
| Scotland | 3,935 | 79.5% | 89.0% | 79.9% | 75.2% | 0% | 92.1% | 0% | 0% |
| Wales | 2,358 | 63.1% | 90.2% | 44.5% | 42.9% | 70.9% | 71.5% | 33.6% | 59.0% |
| UK | 44,513 | 76.1% | 89.5% | 66.0% | 62.9% | 32.8% | 54.1% | 35.9% | 38.2% |

Table E: Data completeness for invasive breast cancers diagnosed in females in 2007 (expressed as % of surgically treated invasive tumours)

| English region/ Celtic country | Number of invasive tumours | % with known invasive tumour size | % with known invasive tumour grade | % with known nodal status | % with known NPI | % with known vascular invasion status | % with known ER status | % with known PgR status | % with known HER2 status |
|-----------------------------------|----------------------------|-----------------------------------|------------------------------------|---------------------------|------------------|---------------------------------------|------------------------|-------------------------|--------------------------|
| Eastern | 3,532 | 96.3% | 96.9% | 92.8% | 90.6% | 18.7% | 50.1% | 31.3% | 44.0% |
| North West | 3,799 | 93.5% | 96.7% | 89.4% | 87.4% | 14.9% | 52.1% | 51.6% | 43.9% |
| Northern & Yorkshire | 3,959 | 96.5% | 97.9% | 65.3% | 64.7% | 31.9% | 65.6% | 57.8% | 55.8% |
| Oxford | 1,744 | 86.9% | 98.7% | 94.8% | 83.1% | 92.3% | 47.2% | 47.0% | 40.5% |
| South West | 4,847 | 94.4% | 96.5% | 89.9% | 87.8% | 83.7% | 57.9% | 42.7% | 43.4% |
| Thames | 6,057 | 85.1% | 93.8% | 61.5% | 56.2% | 9.7% | 36.7% | 34.4% | 30.4% |
| Trent | 2,726 | 52.7% | 97.1% | 52.1% | 51.0% | 14.5% | 52.4% | 14.2% | 47.7% |
| W Midlands | 3,525 | 98.3% | 98.6% | 96.3% | 94.7% | 93.0% | 99.0% | 89.5% | 91.8% |
| England | 30,189 | 89.2% | 96.6% | 78.9% | 75.9% | 41.1% | 56.7% | 45.9% | 48.4% |
| N Ireland | 882 | 93.3% | 93.0% | 93.7% | 85.0% | 0% | 0% | 0% | 0% |
| Scotland | 3,119 | 94.8% | 97.3% | 95.6% | 90.5% | 0% | 94.1% | 0% | 0% |
| Wales | 1,786 | 79.9% | 96.4% | 57.1% | 55.1% | 85.8% | 78.7% | 37.0% | 67.1% |
| UK | 35,976 | 89.3% | 96.6% | 79.6% | 76.4% | 38.8% | 59.7% | 40.4% | 44.0% |

Appendix 4 - Acronyms

| Abbreviation | Full text |
|---------------------|--|
| ABS | Association of Breast Surgery |
| BCCOM | Breast Cancer Clinical Outcome Measures |
| BCS | Breast Conserving Surgery |
| DCIS | Ductal Carcinoma In Situ |
| DCO | Death Certificate Only |
| EPG | Excellent Prognostic Group |
| ER | Oestrogen Receptor |
| GPG | Good Prognostic Group |
| HER2 | Human Epidermal Growth Factor 2 |
| HES | Hospital Episode Statistics |
| ID2007 | Index of Multiple Deprivation 2007 |
| LSOA | Lower Super Output Area |
| MPG1 | Moderate Prognostic Group 1 |
| MPG2 | Moderate Prognostic Group 2 |
| MX | Mastectomy |
| NBSS | National Breast Screening System |
| NCIN | National Cancer Intelligence Network |
| NHS | National Health Service |
| NHSBSP | National Health Service Breast Screening Programme |
| NIMDM | Northern Ireland Multiple Deprivation Measure 2005 |
| NPI | Nottingham Prognostic Index |
| OA | Output Area |
| ONS | Office for National Statistics |
| PEDW | Patient Episode Database for Wales |
| PPG | Poor Prognostic Group |
| QA | Quality Assurance |
| SIMD | Scottish Index of Multiple Deprivation 2006 |
| SLNB | Sentinel Lymph Node Biopsy |
| SMR01 | Scottish Morbidity Record |
| WIMD | Welsh Index of Multiple Deprivation 2008 |
| WMCIU | West Midlands Cancer Intelligence Unit |

MHS